

Full Length Research Paper

Assessment of Microcystin Presence in Freshwater Ecosystems of Southern Mozambique

Olivia Pedro^{1,2,4*}, Thomas Rundberget³, Elisabeth Lie¹, Dacia Correia^{2,4}
Janneche U. Skaare^{1,3}, Knut G. Berdal³, Luis Neves^{2,4}, Morten Sandvik³

¹Norwegian School of Veterinary Science, P.B. 8146 Dep, NO-0033 Oslo, Norway ²Veterinary Faculty of the Eduardo Mondlane University, Av. De Mozambique km 1.5, P.B. 257 Maputo, Mozambique ³Norwegian Veterinary Institute, Ullevålsveien 68, P.B. 750 Sentrum, NO-0106 Oslo, Norway ⁴Centro de Biotecnologia da Universidade Eduardo Mondlane, Av. De Mozambique km 1.5, P.B. 257 Maputo, Mozambique.

Accepted 02 February, 2024

Many countries have started to monitor cyanobacterial cell densities and microcystin concentrations in raw water sources and recreational water due to human and animal poisonings and the results of toxicological studies, which have shown the adverse effects of microcystins to some mammals. Mozambique has no reports of intoxications or deaths of animals or human by cyanobacterial toxins. However, blooms of cyanobacteria have been reported in some aquatic ecosystems. The aim of this study was to detect and quantify the microcystins extracted in water samples from Pequenos Libombos Dam, Nhambavale Lake and Chòkwé Irrigation Channel, all located in southern Mozambique using LC-MS. The Microcystins MC-LR, -YR and -RR were detected in the three sampling areas. MC-LR was found in higher concentrations than the other two variants. The highest total concentration of microcystins detected was 7.89 µg/L in water samples from Nhambavale Lake. Pequenos Libombos Dam revealed concentrations of MC-LR at below quantifiable levels. Based on these results, it is recommended that drinking water supplies in Mozambique are monitored and that the risks of human and animal intoxication by cyanotoxins are assessed and managed.

Keywords: Mozambique, LC-MS, MC-LR, Microcystins.

INTRODUCTION

Mozambique is a developing country located in the southeast coast of Africa (10° 30'-26° 52'S and 40° 50'-30° - 31' E). The country covers a total land area of 800,000 km² and has a wide range of freshwater bodies used as sources of drinking water. Few studies have

Abbreviations: MC, Microcystin; PSDs, Passive sampling devices; LC-MS, Liquid chromatography-mass spectrometry; ELISA, Enzyme linked immunosorbent assay; PL, Pequenos Libombos Dam; NL, Nhambavale lake; CH, Chòkwé Irrigation Channel.

focused on cyanobacteria and their toxin in Mozambique (Pedro *et al.* 2011; Mussagy *et al.* 2006; Bojcevska and Jergil 2003). Given the limited resources of the country, problems of aquatic pollution have not received the proper attention. Toxic cyanobacterial blooms are commonly found in fresh, brackish and marine water. A survey in freshwater has shown that in average, approximately 60% of the blooms in freshwater contain toxins of cyanobacteria (Sivonen 2007). The most common cyanobacterial toxins (cyanotoxins) are hepatotoxins (microcystins and nodularins), neurotoxins (anatoxin-a, anatoxin-a(S) and saxitoxins) and citotoxins (cylindrospermopsins) (Sivonen 2007). Microcystins are the most prevalent cyanobacterial hepatotoxins in freshwater and are most frequently produced by strains of the genera *Anabaena*, *Microcystis*, and *Plan-ktothrix* (formerly *Oscillatoria*) (Sivonen and Jones 1999).

*Corresponding Author E-mail: oliviapedro@uem.mz
olypedro76@gmail.com

Moreover, the involvement of *Anabaenopsis*, *Hapalosiphon*, and *Nostoc* have been less frequently reported (Dittmann and Börner 2005; Vaitomaa *et al.* 2003; Lyra *et al.* 2001; Hummert *et al.* 2001). The microcystins are composed of five common amino acids and pairs of L-amino acids as variants. The most common ones are methyl aspartic acid, alanine, N-methyldehydro-alanine, glutamic acid, and a unique amino acid called Adda (3-amino-9-methoxy-2,6,8-trimethyl-10-phenyldeca-4,6-dienoic acid) (Takino and Kyono 2000).

More than 80 different analogues of microcystins have been described from natural blooms or laboratory cultures of cyanobacteria (Krüger *et al.* 2009; Welker and van Döhren 2006), and most of these differ in the structures of the amino acid residues in positions 2, 3, 4 and 7 of the cyclic peptides (Eckart 1994). Microcystins contain N-methyldehydroalanine (Mdha) or Dehydroalanine (Dha) at unit 7 (Kaya *et al.* 2001). MC-LR as well as all other variants of microcystins has tumor promotion activity (Nishiwaki-Matsushima *et al.* 1992), however there is no evidence of cancer-related activities in the case of [Dhb⁷] microcystins (Kaya *et al.* 2001).

Microcystins inhibit eukaryotic serine/threonine protein phosphatase 1 and 2A, resulting in excess phosphorylation of cytoskeletal filaments. This may cause massive hepatic hemorrhage and death (Znachor *et al.* 2006; Runnegar *et al.* 1991; Romanowska-Duda *et al.* 2002; Honkanen *et al.* 1990; MacKintosh *et al.* 1990; Carmichael 1994; Vaitomaa *et al.* 2003). Although there are several potential routes of exposure for microcystins (oral consumption, inhalation, or skin absorption), the most common is ingestion of water (Znachor *et al.* 2006). It is also noteworthy that microcystins are very stable and may accumulate in the food chain (Watanabe *et al.* 1992). A wide range of wild animals and birds have been affected, with consequences ranging from non-fatal (inhibition of invertebrate feeding, delayed fish egg-hatching) to fatal (mass mortalities of birds, wild and domestic animals and humans) (Metcalf and Codd 2004; Bourke and Hawes 1983; Skulberg *et al.* 1984; Carmichael and Falconer 1993; Ransom *et al.* 1994; Jochimsen *et al.* 1998).

For effective management of cyanobacterial hazards to human health, it is necessary with a basic understanding of the properties, the behaviour in natural ecosystems and the environmental conditions which support the growth of certain species (WHO 1999). Cyanobacterial growth in water bodies is stimulated by nitrogen- and phosphorus-containing nutrients, warm temperatures, stagnant water and intense sunlight (Gobler *et al.* 2007). A high cyanobacteria biomass contributes to aesthetic problems, impairment of recreational use (because of scum and unpleasant odor), affects the taste of treated drinking water and more importantly it poses a serious health risk (Znachor *et al.* 2006; Rantala *et al.* 2006; Sivonen and Jones 1999; Pan

et al. 2002). Identification and quantification of cyanobacteria in water resources is the principal component of cyanotoxin monitoring programmes and can provide an effective early warning system for the development of potentially toxic blooms. The World Health Organization (WHO) has set a provisional guideline of 1 µg/L of total microcystins in drinking water (WHO 2008) and for recreational water there are provisional guidelines of 20000 cells/ml and 100000 cells/ml resulting in low or moderate probabilities of adverse health effects, respectively (Falconer *et al.* 1999; Foulds *et al.* 2002). Ueno *et al.* (1996) have advocated for even stricter guideline value for microcystin LR of 0.1 µg/L due to the potential liver damage since rural communities may be consuming surface water containing cyanotoxins at low level over a period of time. For the risk assessment studies, it is important not only to know the contents of total microcystins but also the presence and relative proportion of each of the microcystin analogue in water collections. (Codd *et al.* 1999).

Many countries in Africa have reported cases of intoxication and deaths of animal that may have been caused by cyanobacterial toxins. In South Africa, *Microcystis aeruginosa* was reported as the most common species of cyanobacteria causing death of livestock (Oberholster *et al.* 2004). Frequent mass mortalities of lesser flamingos have been reported in Kenya Rift Valley (Ballot *et al.* 2004; Ndeti and Munhandiki 2005) and Tanzania (Nonga *et al.* 2011). Cyanobacterial toxicosis was considered the most probable cause of mass mortalities based on the evidence of traces of toxin in the liver of the flamingos. In 1984 massive fish mortality was observed in the Nyanza Gulf of Lake Victoria in Kenya. This fish mortality coincided with the occurrence of cyanobacteria (Ochumba 1990).

Mozambique has no reports of intoxication or mortality on animal or human populations due to cyanobacterial toxins. However, species of cyanobacteria and microcystins have been reported in some aquatic ecosystems by Polymerase Chain reaction (PCR), microscopic observation, Liquid Chromatography coupled to Mass Spectrometry (LC-MS) and Enzyme Linked Immunosorbent Assay (ELISA) (Pedro *et al.* 2011; Mussagy *et al.* 2006; Bojcevska and Jergil 2003). The aim of this study was to investigate the presence of microcystins in water samples collected in the selected water bodies in southern Mozambique using approach LC-MS.

MATERIAL AND METHODS

Study area

This study was carried out in three different sampling areas located in south of Mozambique: Pequenos




Figure 1. Map of south of Mozambique showing the three study areas: Pequenos Libombos Dam, Chokwe Irrigation Channels and Nhambavale Lake. The numbers 1 to 5 indicate the sampling stations at each area.

Pequenos Libombos dam (PL), Nhambavale lake (NL) and Chokwe Irrigation Channel (CH). Pequenos Libombos Dam is a manmade impoundment located 35 km west of Maputo and is the main drinking-water supplier to Maputo, the capital of Mozambique, and also used for fishing. Nhambavale Lake is located in North of Gaza province in Chidenguele region and is also used for drinking, fishing and recreation. A Chokwe irrigation Channel, also located in Gaza Province, is used mainly as source of water for irrigation, but also as a source of drinking water. Water samples were collected in three to five sampling stations at each sampling area (Figure 1). The difference in number of sampling stations was due to the limited accessibility of the sampling areas. The samples used here have been previously analyzed by PCR (Pedro *et al.* 2011).

Cyanobacterial strains and sampling

Sampling took place on 7th of June 2008 and 5th of March 2009. Thirteen water samples were collected during each sampling period (5 samples from Chokwe Irrigation

Channel, 5 from Nhambavale Lake and 3 from Pequenos Libombos dam). In 2008 water samples were collected directly into 1 liter bottles, submersed to about 1 meter below the surface without an additional filtration. In 2009, 30 liters of water from the lake were filtered by conical plankton net (20 μ m mesh) to 500 ml bottles. Samples were stored at -20°C until processing. *Anabaena circinalis* (NIVA-CYA82) and *Anabaena lemmermannii* var. *minor* (NIVA-CYA 83/1), both purchased at NIVA (Norwegian Institute for Water Research) were included as positive and negative control respectively.

Sample preparation for microcystins detection

Extraction of toxins from water samples

Ten ml of water samples and 2 ml of cell cultures were sonicated in an ultrasonic bath for 10 min. Microcystins were extracted from twenty three samples with methanol. Samples were passed through a Bond Elut C₁₈ cartridge (Sample preparation sample, VARIAN^{VA}) previously activated by 1 ml of methanol and washed by distilled

Table 1. Microcystin variants and concentration in freshwater bodies in Mozambique. (PL) Pequenos Libombos Dam, (CH) Chòkwé irrigation channel, (NL) Nhambavale Lake. (08) water samples collected in 2008, (09) water samples collected in 2009. The numbers 1 to 5 indicate the sampling stations.

Sampling area	Samples name	Concentrations (µg/L)			
		MC-LR	MC-YR	MC-RR	Total MC
Pequenos Libombos Dam	1PL-09	bql	-	-	-
	2PL-09	bql	0.10	-	0.10
Chòkwé irrigation channel	4CH-09	0.67	0.06	-	0.73
	5CH-09	0.68	-	-	0.68
Nhambavale Lake	4NL-08	1.65	0.04	-	1.69
	5NL-08	2.6	-	bql	2.6
	3NL-09	2.17	-	-	2.17
	4NL-09	7.82	0.07	-	7.89
	5NL-09	0.86	-	-	0.86

(-) negative samples on the test.
bql- below quantification levels.

Table 2. Microcystins in freshwater bodies in Mozambique in 2008 and 2009.

Sampling area	No. of samples	MC positive samples (%)	MC conc in water (µg/L)	MC variants	pH
Pequenos Libombos Dam	6	2 (33.3)	0.10	LR; YR	8.55 – 8.65
Chòkwé irrigation channel	7	2 (28.5)	0.68 – 0.73	LR; YR	8.20 – 8.56
Nhambavale Lake	10	5 (50)	0.86 – 7.89	LR; YR; RR	8.18 – 8.40

water. Toxins were eluted from cartridge with 2 ml of methanol.

LC-MS analysis.

For analysis of the extracted toxin for detection of microcystins a modified method described by Neffling *et al.* (2010) was used. Liquid chromatography was performed on a XBridge C₁₈ column (3.5 µm, 50 × 2.1 mm) (Waters, Milford, MA, USA), using a Surveyor HPLC system (Thermo Electron Corporation, Waltham, MA, USA). Separation was achieved using linear gradient elution at 0.3 mL/min starting with MeCN–water (30:70, both containing 0.1% formic acid) rising to 100% MeCN over 10 min. Isocratic elution with 100% MeCN was maintained for 5 min before the eluent was switched back to 30% MeCN. The HPLC system was coupled to an LTQ ion trap mass spectrometer operating with an electrospray ionization (ESI) interface (Thermo Electron Corporation, Waltham, MA, USA). Typical ESI parameters were spray voltage 4 kV, heated capillary temperature 250 °C and sheath gas 60 units (ca 60 L/h) of N₂. The mass spectrometer was operated in scan mode (*m/z* 400–1300) and all samples were diluted to appropriate concentrations with 60% MeOH. For full scan MS/MS analyses, the selected precursor ions were isolated with an isolation width of 3 Da. The detection limit of the method was 9 ng/ml. Microcystins -LR, -YR

and -RR were included as standards in the study. LC-MS was conducted at Norwegian Veterinary Institute (Norway).

RESULTS AND DISCUSSION

LC-MS analysis revealed the presence of three variants of microcystins in water samples collected in Southern Mozambique (Table 1 and 2). MC-LR, -YR, and -RR were measured in the samples on the basis of both their mass spectra and retention time. Peaks at 995.6 *m/z* ([M+H]⁺), 1045.5 *m/z* ([M+H]⁺) and 1038.5 *m/z* ([M+2H]²⁺) and retention times of 3.87, 3.72 and 3.07 min correspondent to MC- LR, -YR and -RR respectively were identified in some water samples (Table 1, Figure 2) as well as in positive control strain *Anabaena circinalis* (NIVA-CYA 82). Table 2 shows the summary of the concentration of microcystins in water samples collected in Nhambavale Lake, Chòkwé Irrigation Channel and Pequenos Libombos dam in June 7th 2008 and March 5th 2009. Microcystins were detected in 7 out of the 23 water samples analyzed (Table 2), with total concentration ranging from 0.10 µg/L to 7.89 µg/L. The highest concentration of total microcystins (7.89 µg/L) was detected in Nhambavale lake (Table 2) and was around 10 times higher compared with the microcystin concentrations measured in Chòkwé Irrigation Channel (the highest concentration was 0.73 µg/L) and around 7




Figure 2. LC–MS chromatogram and MS² spectra of microcystins LR and YR of water sample collected in Nhambavale Lake, Mozambique.

Times above the WHO guideline of 1 µg/L total microcystins in drinking water (WHO 2008).

Figure 3 shows that the LR variant was found to be the predominant microcystin in the all three sampling areas. The concentration of MC-YR varied from 0.04 µg/L to 0.1 µg/L of water and was detected in only 4 out of 23 water samples analyzed. MC-RR was detected only in Nhambavale Lake and was at below quantifiable levels. MC-LR was also detected at below quantifiable level at Pequenos Libombos Dam (Table 1). High concentration of microcystins was observed in March 2009, a period characterized by high temperatures (21–26°C). Warm temperatures are normally associated with high concentrations of microcystins (Neilan *et al.* 1994; Wicks and Thiel 1990; Chorus and Bartram 1999; Jöhnk *et al.* 2008; Paerl and Huisman 2008). Additionally, different factors such as sampling collection method and environmental conditions in the two years may be contributed to the differences in the microcystin concentrations measured.

An ELISA assay has previously demonstrated the presence of microcystins in Nhambavale Lake, with concentration ranging from 0.5 to 6.8 µg/L (Bojcevska and Jergil 2003) which were almost the same as this study (7.89 µg/L) and 7 times above the WHO guideline

of 1 µg/L total microcystins (WHO 2008). Previous studies in Chôkwé, have indicated concentration of microcystins below 0.1 µg/L (Bojcevska and Jergil 2003). In the present study MC-LR varied from 0.67 to 0.68 µg/L, indicating that microcystin concentration levels in this area usually may stay below the WHO threshold. Analysis for microcystins in passive sampling devices (PSDs) by LC-MS revealed 3 microcystin variants in Southern Mozambique (MC-LR, -YR and -RR) at the concentrations of 2.1– 159.4 ng/g of PSDs (Pedro *et al.* 2011). During the sampling no bloom formation was observed in the three areas, however Bojcevska and Jergil (2003) reported bloom formation in Pequenos Libombos Dam. This lack of consistent information strongly points out the urgent need for longitudinal screening studies in all relevant areas of Mozambique. ELISA method used previously to quantify microcystins in Mozambique is a good screening test, however it has the limitation of being expensive and not differentiates between variants of microcystins as the LC-MS method does.

Other studies in Africa reported high concentration of microcystins, with values above the WHO threshold. Microcystin concentrations ranging from 2 µg/L to 23.7 µg/L have been reported from different water bodies in




Figure 3. Microcystin concentrations (mean) in samples collected in 2008 (Nhambavale Lake) and 2009 (Pequenos Libombos Dam, Chòkwé Irrigation Schemes, and Nhambavale Lake). Pequenos Libombos Dam in 2009 showed MC-LR concentration at below quantifiable levels.

Southern Africa (Makhera *et al.* 2011; Oberholster *et al.* 2009). Okello *et al.* (2011) reported concentration of microcystins varying from 0.02 µg/L to 10 µg/L of water in Uganda. Microcystin concentrations found in the present study are within the range detected in different studies in Africa, which indicate the great variability of this parameter. Furthermore, microcystin concentrations are likely to vary substantially within the same water collection during the year and over a period of several years. Drinking water may be the major source of microcystins at the present time, based on the water intake of people every day for a long time. However, the possibility of exposure to microcystins through the food chain cannot be excluded (Ueno *et al.* 1996). Programs for monitoring water bodies should be considered by the responsible authorities in Mozambique.

The concentration of microcystins in water samples from the three areas varied from non-detectable levels in Pequenos Libombos dam to high levels (7.89 µg/L) and above the WHO guideline of total microcystins in water in Nhambavale Lake. Nhambavale Lake is a touristic area and populations from Chidenguele Village take untreated water for consumption and do hand washing into the lake. The lake is also used to practice recreation activities such as sailing, canoeing and fishing. Our results show that populations from Nhambavale Lake may be consuming surface water containing cyanotoxins that potentially could increase the risk of liver cancer. Carmichael (1994) and Juković *et al.* (2008) found that

the incidence of primary liver cancer is higher in peoples who commonly use pond or river water for consumption. Ueno *et al.* (1996) also suggest that the consumption of microcystins at low level for long term may result also in liver damage. Conversely, Pequenos Libombos Dam is a source of water which supply Maputo city, after water treatment. However, rural populations have access to untreated water from the Incomati River that supply this dam. The conventional methods for water treatment are not effective for elimination of microcystins. Volterra *et al.* (1990) reported that trichomes of cyanobacteria may be disrupted during drinking water treatment, and their hepatotoxins released into the drinking water (Himberg *et al.* 1989; Brittain *et al.* 2000).

In conclusion, the presence of hepatotoxic microcystins in freshwater from Mozambique may constitute a health risk especially for rural communities that may be in contact with the contaminated water without any form of treatment. Taking into consideration the highest concentration of microcystins detected in the lakes (7.89 µg/L), it is important that drinking water supplies in Mozambique are monitored for the presence of microcystins to manage the risk of intoxication.

ACKNOWLEDGEMENTS

This study was supported by NUFU Program: ZOOTOX-2007-2011. We are grateful to Norwegian Veterinary

Institute and Laboratory of Environmental Toxicology at the Norwegian School of Veterinary Science for allowing use of the laboratory and all the required equipment. We thanks the administration of ARA-SUL (Pequenos Libombos), of the Village of Chidenguele, and HICEP (Hidráulica de Chokwé Empresa Pública), for allowing sampling collection. All Staff of Biotechnology Center in Mozambique, especially Joelma Leão, Paula Macucule and Iris Victorino, are personally thanked for their assistance in the sampling process.

REFERENCES

- Ballot A, Krienitz L, Kotut K, Wiegand C, Metcalf JS, Codd GA, Pflugmacher S (2004). Cyanobacteria and Cyanobacterial Toxins in three Alkaline Rift Valley Lakes of Kenya—Lakes Bogoria, Nakuru and Elmenteita. *J. Plank. Res.* 26 (8): 925–935.
- Biol. 14, 495–512.
- Bojcevska H, Jergil E (2003). Master Degree Projec. Removal of cyanobacterial toxins (LPS endotoxin and microcystin) in drinking-water using the BioSand household water filter. Upsala University, Sweden, pp 1-44.
- Bourke ATC, Hawes RB (1983). Fresh water cyanobacteria (blue-green algae) and human health. *Med. J. Aust.* 1:491-492.
- Brittain SM, Wang J, Babcock-Jackson L, Carmichael WW, Rinehart KL, Culver DA (2000). Isolation and characterization of microcystins, cyclic heptapeptide hepatotoxins from a Lake Erie Strain of *Microcystis aeruginosa*. *J. Great. Lakes. Res.* 26:241–249.
- Carmichael WW (1994). The toxins of cyanobacteria. *Sci. Am.* 270:64-72.
- Carmichael WW, Falconer IR (1993). Diseases related to freshwater blue-green algal toxin, and control measures. In *Algal Toxins in Seafood and Drinking Water* (Farconer, I.R. Ed). Pp 187-209, Academic press, New York.
- Chorus I, Bartram J (1999). *Toxic Cyanobacteria in Water. A Guide to Their Public Health Consequences, Monitoring and Management.* E & FN Spon, London.
- Codd GA, Bell SG, Kaya K, Ward CJ, Beattie KA, Metcalf JS (1999). Cyanobacterial toxins, exposure routes and human health. *Eur. J. Phycol.* 34, 405-415.
- Dittman E, Börner T (2005). Genetic contributions to the risk assessment of microcystin in the environment. *Toxicol. Appl. Pharmacol.* 3: 192–200.
- Eckart K (1994). Mass spectrometry of cyclic peptides. *Mass. Spectrom. Rev.* 13: 23-55.
- Falconer I, Bartram J, Chorus I, Kuiper- Goodman T, Utkilen H, Burch M, Codd GA (1999). Safe levels and safe practices. P. 155-178. In Chorus, I., Bartram, J. (ed.). *Toxic cyanobacteria in water – a guide to their public health consequences, monitoring and management.* E. & F.N. Spon. London.
- Foulds IV, Granacki A, Xiao C, Krull UJ, Castle A, Horgen PA (2002). Quantification of microcystin-producing cyanobacteria and *E. coli* in water by 5- nuclease PCR. *J. Appl. Microbiol.* 93:825–834.
- Gobler CG, Davis TW, Coyne KJ, Boyer GL (2007). Interactive influences of nutrient loading, zooplankton grazing, and microcystin synthetase gene expression on cyanobacterial bloom dynamics in a eutrophic New York lake. *Harm. Algae.* 6: 119–133.
- Himberg K, Keijola AM, Hiisvirta L, Pyysalo H, Sivonen K (1989). The effect of water treatment processes on the removal of hepatotoxins from *Microcystis* and *Oscillatoria* cyanobacteria: a laboratory study. *Wat. Res.* 23(8): 979–984.
- Honkanen RE, Zwiller J, Moore RE, Daily SL, Khatra BS, Dukelow M, Boynton AL (1990). Characterization of microcystin-LR, a potent inhibitor of type 1 and 2A protein phosphatases. *J. Biol. Chem.* 265:19401-19404.
- Hummert C, Dahlmann J, Reichel M, Luckas B (2001). Analytical techniques for monitoring harmful cyanobacteria in lakes. *Lakes Reser. Res.Manage.* 6: 159–168.
- Jochimsen EM, Carmichael WW, An J, Cardo DM, Cookson ST, Holmes CEM, de C Antunes MB, de Melo Filho DA, Lyra TM, Barreto VST, Azevedo SMFO, Jarvis WR (1998). Liver failure and death after exposure to microcystins at a hemodialysis center in Brazil. *N. Engl. J. Med.* 338:873-878.
- Jöhnk KD, Huisman J, Sharples J, Sommeijer B, Visser PM, Strooms JM (2008). Summer heatwaves promote blooms of harmful cyanobacteria. *Global Change*
- Juković MJ, Svirčev ZB, Baltić VV, Stojanović DB, Mihajlović AI, Baltić MV (2008). Cyanotoxins – new health risk factor in Serbia. *Review Article. Arch. Oncol.* 16(3-4): 55-58.
- Kaya K, Sano T, Inove H, Takagi HA (2001). Selective determination of total normal microcystin by colorimetry: LC/UV detection and/or LC/MS. *Anal. Chim. Acta.* 450: 73-80.
- Krüger T, Christian B, Luckas B (2009). Development of an analytical method for the unambiguous structure elucidation of cyclic peptides with special appliance for hepatotoxic desmethylated microcystins. *Toxicon* 54 (3): 302–312.
- Lyra C, Suomalainen S, Gugger M, Vezie C, Sundman P, Paulin L, Sivonen K (2001). Molecular characterization of planktic cyanobacteria of *Anabaena*, *Aphanizomenon*, *Microcystis* and *Planktothrix* genera. *Int. J. Syst. Evol. Microbiol.* 51: 513– 526.
- MacKintosh C, Beattie KA, Klumpp S, Cohen P, Codd GA (1990). Cyanobacterial microcystin-LR is a potent and specific inhibitor of protein phosphatases 1 and 2A from both mammals and higher plants. *FEBS.* 264:187-192.
- Makhera M, Gumbo JR, Chigayo K (2011). Monitoring of microcystin LR in Luvuvhu River catchment: Implications for human health. *Afri. J. Biotechnol.* 10 (3): 405-412.
- Metcalf JS, Codd GA (2004). Cyanobacterial toxin in water environment: A review of current knowledge. *Foundation for Water Research, Marlow, UK*, pp 1-36.
- Mussagy A, Annadotter H, Cronberg G (2006). An experimental study of toxin production in *Arthrospira fusiformis* (Cyanophyceae) isolated from African waters. *Toxicon.* 48: 1027–1034.
- Ndetei R, Muhandiki VS (2005). Mortalities of lesser flamingos in Kenyan Rift Valley saline lakes and the implications for sustainable management of the lakes. *Lakes Reserv. Res. Manage.* 10: 51–58.
- Neffling M-R, Lance E, Meriluoto J (2010). Detection of free and covalently bound microcystins in animal tissues by liquid chromatography–tandem mass spectrometry. *Environmental Pollution* 158: 948–952.
- Neilan BA, Hawkins PR, Cox PT, Goodman AE (1994). Towards a molecular taxonomy for the bloom-forming cyano- bacteria. *Aust. J. Mar. Freshwater Res.* 45: 869-873.
- Nishiwaki-Matsushima R, Ohta T, Nishiwaki S, Suganuma M, Kohyama K, Ishikawa T, Carmichael WW, Fujiki H (1992). Liver tumor promotion by the cyanobacterial cyclic peptide toxin microcystin-LR. *J. Cancer. Res. Clin. Oncol.* 118:420-424.
- Nonga HE, Sandvik M, Miles CO, Lie E, Mdegela RH, Mwamengele GL, Semuguruka WD, Skaare JU (2011). Possible involvement of microcystins in the unexplained mass mortalities of Lesser Flamingo (*Phoeniconaias minor* Geoffroy) at Lake Manyara in Tanzania. *Hydrobiologia.* DOI 10.1007/s10750-011-0844-8.
- Oberholster PJ, Botha A-M, Grobbelaar JU (2004). *Microcystis aeruginosa*: source of toxic microcystins in drinking water Review. *Afri. J. Biotechnol.* 3 (3): 159-168.
- Oberholster PJ, Myburgh JG, Govender D, Bengis R, Botha A-M (2009). Identification of toxigenic *Microcystis* strains after incidents of wild animals mortalities in the Kruger National Park, South Africa. *Ecotox. Environ. Saf.* 72: 1177- 1182.
- Ochumba PBO (1990). Massive fish kills within the Nyanza gulf of lake Victoria, Kenya. *Hydrobiologia.* 208(1/2):93-99.
- Okello O and Kurmayer R (2011). Lakes & Reservoirs Seasonal development of cyanobacteria and microcystin production in Ugandan freshwater lakes. *Res. Manage.* 16: 123–135.
- Paerl HW, Huisman J (2008). Blooms like it hot. *Science* 320, 57–58.
- Pan H, Song L, Liu Y, Börner T (2002). Detection of hepatotoxic *Microcystis* strains by PCR with intact cells from both culture and environmental samples. *Arch. Microbiol.* 178: 421–427.

- Pedro O, Correia D, Lie E, Skåre JU, Leão J, Neves L, Sandvik M, Berdal KG (2011). Polymerase chain reaction (PCR) detection of the predominant microcystin-producing genotype of cyanobacteria in Mozambican lakes. *Afri. J. Biotechnol.* 10(83): 19299-19308.
- Rantala A, Rajaniemi-Wacklin P, Lyra C, Lepisto L, Rintala J, Mankiewicz-Boczek J, Sivonen K (2006). Detection of Microcystin-Producing Cyanobacteria in Finnish Lakes with Genus-Specific Microcystin Synthetase Gene E (*mcyE*) PCR and Associations with Environmental Factors. *Appl. Environ. Microbiol.* 72 (9): 6101–6110.
- Ressom R, Soong FS, Fitzgerald J, Turczynowicz L, El Saadi O, Roder D, Maynard T, Falconer I (1994). Health effects of toxic cyanobacteria (blue-green algae). National Health and Medical Research Council, Australian Government Publishing Service, Canberra, Australia, pp 108.
- Romanowska-Duda Z, Mankiewicz J, Tarczynska M, Walter Z, Zalewski M (2002). The effect of toxic cyanobacteria (blue-green algae) on water plants and animal cells. *Polish. J. Environ. Studies.* 11: 561-566.
- Runnegar MT, Gerdes RG, Falconer IR (1991). The uptake of the cyanobacterial epatoxins microcystins by isolated rat hepatocytes. *Toxicon.* 29: 43-51.
- Sivonen K (2007). Emerging high throughput analyses of cyanobacterial toxins and toxic cyanobacteria. *Proceedings of the Interagency, International Symposium on Cyanobacterial Harmful Alga Blooms.* *Adv. Experim. Med. Biol.* 523-542.
- Sivonen K, Jones G (1999). Cyanobacterial toxins, p. 41–111. In I. Chorus and J. Bartram (ed.), *Toxic cyanobacteria in water. A guide to their public health consequences, monitoring and management.* E and FN Spon, London, United Kingdom.
- Skulberg OM, Codd GA, Carmichael WW (1984). Toxic blue-green algal bloom in Europe: A growing problem. *Anbio.* 13: 244-247.
- Takino M, Kyono Y (2000). LC/MS Analysis of Microcystins in Freshwater by Electrospray Ionization. *Application Brief.* Agilent Technologies, Inc. USA.
- Ueno Y, Nagata S, Tsutsumi T, Hasegawa A, Watanabe MF, Park HD, Chen GE, Yu SH (1996). Detection of microcystins, a blue-green algal hepatotoxin, in drinking water sampled in Haimen and Fusui, endemic areas of primary liver cancer in China, by highly sensitive immunoassay. *Carcinogenesis.* 17 (6): 1317- 1321.
- Vaitomaa J, Rantala A, Halinen K, Rouhiainen L, Tallberg P, Møkelke L, Sivonen K (2003). Quantitative real-time PCR for determination of microcystin synthetase *e* copy numbers for *Microcystis* and *Anabaena* in lakes. *Appl. Environ. Microbiol.* 69: 7289– 7297.
- Volterra L, Bruno M, Gucci PM, Pierdominici E (1990). Cyanobacteria monitoring and health related problems. In: Castilio, G., Campos, V., Herrera, L. (Eds.), *Proceedings of the second Biennial Water Quality Symposium: Microbiological aspects* Vina del Mar. University of Chile Press, Vina del Mar, Chile, pp. 25-30.
- Watanabe MF, Tsuji K, Watanabe Y, Harada K-I, Suzuki M (1992). Release of heptapeptide toxin (microcystin) during the decomposition process of *Microcystis aeruginosa*. *Na. Toxins.* 1: 48-53.
- Welker M, Von Dohren H (2006). Cyanobacterial peptides – nature's own combinatorial biosynthesis, *FEMS. Microbiol. Rev.* 30: 530-63.
- WHO, World Health Organization (1998). *Toxic Cyanobacteria in Water: A guide to their public health consequences, monitoring and management.* British Library Cataloguing.
- WHO, World Health Organization (2008). *Guidelines for Drinking Water Quality. (3rd edn incorporating the first and second ADDENDA).* Volume 1. WHO Health Organization Library cataloguing. Geneva, Witzerland.
- Wicks RJ, Thiel PG (1990). Environmental factors affecting the production of peptide toxins in floating scums of cyanobacterium *Microcystis aeruginosa* in a hypertrophic African reservoir. *Environ. Sci. Technol.* 24, 1413-1418.
- Znachor P, Jurczak T, Komárkova J, Jezberová J, Mankiewicz J, Kaštovská K, Zapomělová E (2006). Summer changes in cyanobacterial bloom composition and microcystin concentration in eutrophic Czech Reservoirs. *Environ. Toxicol.* 236- 243.