

## Full Length Research Paper

# Prevalence and risk factors of intestinal parasites among Delgi school children, North Gondar, Ethiopia

Asrat Ayalew<sup>1\*</sup>, Tewodros Debebe<sup>2</sup> and Alemayehu Worku<sup>3</sup><sup>1</sup>Center of Competence, Amhara Regional State Education Agency, P. O. Box 572 Bahir Dar, Ethiopia.<sup>2</sup>Department of Microbiology, Immunology and Parasitology, College of Medicine and Health Sciences, Bahir Dar University, P.O. Box 79 Bahir Dar, Ethiopia.<sup>3</sup>Department of Public Health, Addis Ababa University P. O. Box 1176 Addis Ababa, Ethiopia.

Accepted 06 March, 2019

Epidemiological information on the prevalence of various intestinal parasitic infections and identification of local risk factors in different regions/localities is a prerequisite to develop appropriate control strategies. Hence the aim of this study was to determine the prevalence of intestinal parasites and risk factors among school children. A cross sectional study involving 704 school children was conducted from 27 September to 6 December, 2010. Structured questionnaire were used to identify environmental, socio demographic and behavioral factors. Stool specimens were collected from all study subjects and examined for parasites using direct smear and formol-ether concentration technique. Finally, data entry and analysis was done using Epi-info and SPSS statistical soft ware respectively. Ten species of intestinal parasites were identified with an overall prevalence of 79.8% (562 of 704 children). The most prevalent intestinal parasites identified were *Ascaris lumbricoides* 338 (48%), *Giardia lamblia* 295 (41.9%), *Entamoeba histolytica/dispar* 192 (27.3%), *Schistosoma mansoni* 112 (15.9%), Hookworm 81(11.5%). Of the total positive stool samples 96 (13.6%) were single infection and 466 (66.2%) were mixed infection, from this mixed infections majority of the children had double infection 309 (43.9%). Intestinal parasite prevalence was higher in children with less educated mother, in children who have habit of eating raw/ unwashed vegetables, drinking unprotected well/spring water and who do not have hand washing practice before meal ( $P < 0.05$ ). In conclusion, Intestinal parasite infection is a health problem among Delgi school children. Therefore, interventions including improvement of sanitation, provision of safe water, and health education on personal hygiene to the students and to their parents should be given.

**Key words:** Intestinal parasites, Delgi school children, Ethiopia.

## INTRODUCTION

Intestinal parasitic infections are among the most common infections worldwide. It is estimated that 3.5 billion people are affected, and 450 million are ill as a result of these infections, the majority being children (WHO, 1998). Epidemiological studies carried out in different countries have shown that the situation of an individual is an important cause in the prevalence of intestinal parasitic infection, having a greater rate in

children (Udonsi and Amabibi, 1992; Nokes et al., 1992). Inadequate water sanitation and hygiene are responsible for a major proportion of the burden of disease and death in developing countries (Merid et al., 2001). In addition, intestinal parasitic agents increase in polluted environments such as refuse heaps, gutters and swage units in and around human dwelling and living conditions of the people in crowded or unhealthy situations (Phiri et al., 2000).

The level of harm caused by intestinal parasite infection to the health of individual and communities depend on: the parasite species, the nature of the interaction between the parasite and the concurrent infections, the

\*Corresponding author: E-mail: [asratayalew@yahoo.com](mailto:asratayalew@yahoo.com). Tel: +251(0)911630599.

intensity and course of infection and nutritional and immunological status of the population (WHO, 1987; Hadidjaja et al., 1998). The common consequences of intestinal parasitic infections have been shown to affect nutritional status, physical development, mental function, verbal ability and cognitive behavior in children (Stephenson et al., 1993; Nokes and Bundy, 1994; Hadidjaja et al., 1998). Similarly, the distribution and prevalence of various species of intestinal parasites also differs from region to region because of several environmental, social and geographical factors. Hence, study on the prevalence of various intestinal parasitic infections is a prerequisite not only for formulation of appropriate control strategies but also to predict risk for communities under consideration.

In Ethiopia intestinal parasitosis is prevalent because of low level of living standards, poor environmental sanitation and ignorance of simple health promoting factors (Wolde-michael et al., 1990; Kloos and Tesfa-Yohannes, 1993; Yeneneh, 1994). But there are still several localities for which epidemiological information is not available including the study area. Therefore, the objective of the present study was to determine the prevalence and risk factors of intestinal parasitic infections among school children in Delgi elementary and junior secondary school.

## **MATERIALS AND METHODS**

### **Study area and population**

This study was conducted in Delgi elementary and Junior Secondary School, Delgi town; located in North-West of Gondar city; North Ethiopia. The town Delgi is located 824 km from the capital city Addis Ababa and surrounded by Lake Tana margin with an altitude of 2100 m above sea level and an average temperature of 25 to 28°C. The area is predominately rural and most residents live in villages as agriculturists growing maize, teff (*Eragrostos tef*) and pepper. The total population of the study area is approximately 16319 with the majority Amahara ethnic group (DMR, 2009).

### **Study design and sampling procedures**

Cross-sectional study design was conducted from 27 September to 6 December, 2010 at Delgi town. The sample size was determined using the single proportion population formula, since the overall prevalence rate (P) of intestinal parasite is not known in the study area; (p) was taken as 50%. For the calculation a 95% confidence interval (z) and a 5% margin of error (d) and design effect of 2 was used to allow for cluster sampling. The calculated study population size was 704 including ten percent of the sample size to minimize errors arising from the likely hood of non compliance. Cluster sampling was used in the selection of the study sample. From 1888 students in the school (grade 1 to grade 8), there were a total number of thirty sections; grade 1 to 4 two sections each, grade 6 and 8 five sections each and grade 5 and 7 six sections each, with an average number of 60 students in each section. From these total thirty sections, twelve clusters (sections) were selected randomly to get the required amount of sample size (n=704), accordingly from

grade 1 to 4 one section each and from grade 5 to 8 two sections each were totally included in the study.

### **Data collection procedure and specimen analysis**

A pre-tested questionnaire based on known risk factors was developed and modified (The questions are listed on Table 2). Data were collected by laboratory technicians who were selected and trained for the purpose. To ensure reliable information, the children were interviewed in their mother tongues. The interview included information on socio demographic data, environmental and behavioral factors. At the time of conversation, interviewers also inspected whether the fingernails of the students were trimmed and their foot wear. After checking the completion of the questioners, proper stool samples were taken with labeled clean plastic container, toilet tissue paper, pieces of applicator sticks from all students. As soon as the stool samples presented all specimens were checked for their label, quantity, time, procedure of collection and stool examinations were done by laboratory technologists using direct technique (saline and iodine mounts) and formol-ether concentration technique to identify and detect each stage of intestinal parasites in less than 20 min of stool sample collection. After completion of stool examination, samples were emulsified in a 10% formalin solution and 20% of stool samples were randomly selected and transported to University of Gondar Parasitological laboratory and rechecked blindly to assure the quality control.

### **Data analysis procedures**

Data entry and analysis was done using Epi-info version 3.5.1 and SPSS version 15 computer software respectively. The base line characteristics of the study population were summarized using medians and ranges for continuous variables, simultaneously proportions and frequencies for categorical variables. Internal comparison was made using logistic regression to determine the independent effect of the variables by calculating the strength of the association between infection and risk factors using odds ratio (OR) and 95% confidence interval (CI). Crude and adjusted OR were computed using bivariate and multivariate logistic regression analysis respectively. P value less than 0.05 (5%) was considered statistically significant.


### **Ethical consideration**

The study was approved by the Ethics Committees of Addis Continental Institute of Public Health (ACIPH) and University of Gondar. Participation was fully voluntary and informed written consent was obtained from each study subject. For those children whose age was under fifteen years or unable to understand the purpose of the study written consent was obtained from their family through school director. Finally appropriate treatment was given to those students who were positive for intestinal parasitic infection by local nurses.

## **RESULTS**

### **Prevalence of intestinal parasites**

A total of 704 students took part in the study and were included in the present analysis. Of these, 358 (50.9%) were male and 346 (49.1%) female. The mean age of the



**Figure 1.** Prevalence of each intestinal parasite based on wet mount and formol-ether concentration technique among Delgi school children, September to December 2010.

**Table 1.** Prevalence of single and mixed intestinal parasite infection among Delgi school children, September to December, 2010.

Type of infection	Total frequency n (%)	Most prevalent intestinal parasite species in each type of infection	
		Species	Total frequency n (%)
Single infection	96(13.6)	G.I	44(6.3)
Double infection	309(43.9)	A.I + G.I	99(14.1)
Triple infection	155(22.3)	A.I + S.m + E.m	28(4)
Quadruple infection	2(0.3)	A.I + S.m + E.m + Hw	2(0.3)

G.I\*=*G. lamblia*; A.I\*=*A. lumbricoides*; E.h\*=*E. histolytica/dipar*; H.w\*=Hookworm; S.m\*=*S. mansoni*.

study subject was 12.8 years with a minimum and maximum age of 7 and 25 years respectively. Majority of the study subjects were reside in urban 441(62.6%). Most of the students mothers' were illiterate 395 (56.1%), primary school 195 (27.3%), secondary and above 114 (16.2%). Majority of students were grade 5 to 8, 480 (68.2%).

Microscopic stool sample examination using wet mount and formol-ether concentration technique showed that infections with various intestinal helminthes and protozoan parasites were common in Delgi school children. Out of the 704 school children examined, ten species of intestinal parasites were identified with an overall prevalence of 562 (79.8%). The most prevalent intestinal parasites identified were *A. lumbricoides* 338 (48%), *G. lamblia* 295 (41.9%), *E. histolytica/dipar*

192(27.3%), *S. mansoni* 112(15.9%), Hookworm 81(11.5). The prevalence of *A. lumbricoides* was the highest compared with other parasite species (Figure 1).

Of the total 562 (79.8%) positive individuals 96 (13.6%) were single infection and 466 (66.2%) were mixed infection; from this mixed infection the majority of the students had double infection 309 (43.9%) (Table 1).

### Analysis of risk factors

Among the potential risk factors explored using univariate analysis, mothers'educational level, hand washing practice before meal, habit of eating raw/ unwashed vegetables, water source for drinking, open field defecation and finger nail condition had statistically

**Table 2.** Bivariate logistic regression analysis for factors potentially associated with Intestinal parasite infection among Delgi school children, September to December 2010.

Risk factors	Intestinal parasites		Crude OR (95% CI)
	Total positive n (%)	Total negative n (%)	
<b>Sex</b>			
Male	290(51.6)	68(47.9)	1.16(0.80,1.68)
Female	272(48.4)	74(52.1)	1.00
<b>Age group in years</b>			
5 to 9	127(26.6)	23(16.2)	1.05(0.57,1.93)
10 to 14	298(53.0)	93(65.5)	0.61(0.38,0.98)
15 <sup>+</sup>	137(24.4)	26(18.3)	1.00
<b>Grade level</b>			
1 to 4	183(32.6)	41(28.9)	1.19(0.79,1.78)
5 to 8	379(67.4)	101(71.1)	1.00
<b>Residence</b>			
Urban	358(63.7)	83(58.5)	1.25(0.86,1.82)
Rural	204(36.3)	59(41.5)	1.00
<b>Mother educational status</b>			
Illiterate	325(57.8)	70(49.3)	2.06(1.28,3.31)
Primary school	158(28.1)	37(26.1)	1.89(1.11, 3.23)
Secondary school and above	79(14.1)	35(34.6)	1.00
<b>Hand washing before eating</b>			
Yes	261(46.4)	110(77.5)	1.00
No	301(53.6)	32(22.5)	3.96 (2.59,6.08)
<b>Practice of eating raw/ unwashed vegetables</b>			
Yes			
No	438(77.9)	67(47.2)	3.95 (2.69,5.81)
	124(21.1)	75(52.8)	1.00
<b>Water source for drinking</b>			
Pipe	315(56.0)	110(77.5)	1.00
River	135(24.0)	21(14.8)	2.25(1.35,1.73)
Unprotected well /Spring	112(19.9)	11(7.7%)	3.54( 1.85,6.85)
<b>Open field Defecation</b>			
Yes	97(17.3)	8(5.6)	3.49 (1.65,7.37)
No	465(82.7)	134(94.4)	1.00
<b>Practice of fingernail trim</b>			
Yes	431(76.7)	125(88.0)	1.00
No	131(23.3)	17(12.0)	2.24(1.30,3.85)

significant association to any intestinal parasite infections identified in the study (Table 2).

Multivariate analysis was performed for all variables

that were significantly associated with any intestinal parasite infection from univariate analysis. After adjustment, open field defecation and finger nail

**Table 3.** Bivariate and multivariate logistic regression analysis for factors potentially associated with Intestinal parasite infection among Delgi school children September to December 2010.

Risk factors	Intestinal parasites		Adjusted OR (95% CI)
	Total positive n (%)	Total negative n (%)	
Mother educational status			
Illiterate	325(57.8)	70(49.3)	2.14(1.27,3.63)
Primary school	158(28.1)	37(26.1)	1.85(1.03,3.35)
Secondary school and above	79(14.1)	35(34.6)	1.00
Hand washing before eating			
Yes	261(46.4)	110(77.5)	1.00
No	301(53.6)	32(22.5)	3.88 (2.46,6.08)
Eating practice of raw/ unwashed vegetables			
Yes	438(77.9)	67(47.2)	3.62(2.40,5.45)
No	124(21.1)	75(52.8)	1.00
Water source for drinking			
Pipe	315(56.0)	110(77.5)	1.00
River	135(24.0)	21(14.8)	1.86(1.08,3.20)
Unprotected well/spring	112(19.9)	11(7.7)	3.42(1.71,6.83)

condition were excluded from the model ( $P>0.05$ ). All other variables remained significant positive association with any intestinal parasite infection; these include children who have illiterate mothers (Odds Ratio 2.14; 95% CI 1.27 to 3.63) and primary school (Odds Ratio 1.85; 95%CI 1.03 to 3.35) were more likely to acquire intestinal parasite infection than secondary school and above. In addition, children who did not wash their hands before eating (Odds Ratio 3.88; 95% CI 2.46 to 6.08) were more likely to acquire intestinal parasite infection than children who wash their hands before meal. Similarly, children who had habit of eating raw/unwashed vegetables (Odds Ratio 3.62 95% CI 2.40 to 5.45) were more likely to acquire intestinal parasite infection. Students who drink unprotected well/spring water (Odds Ratio 3.42; 95 CI 1.71 to 6.83) and who drink river water (Odds Ratio; 1.86; 95% CI 1.08 to 3.20) were more likely to acquire intestinal parasite infection as compared to those who drink pipe water (Table 3).

## DISCUSSION

Epidemiological studies carried out in different countries have shown that the situation of an individual is an important cause in the prevalence of intestinal parasitic infection, having a greater rate in children. In line with this view, the present study attempted to assess the prevalence of different intestinal parasitic infections and associated risk factors in school children of Delgi, North

Gondar. In this study, the observed overall prevalence rate of intestinal parasite was found to be (79.8%) relatively higher than previously reported among different regions in Ethiopia, (Jemaneh, 1997; Assefa et al., 1998; Jemaneh, 2000; Tadesse, 2005) and other countries (Magambo et al., 1998; Glickman et al., 1999; Chandrasekhar and Nagesha, 2003; Pinar et al., 2004; Ahsan et al., 2005; Agbolade et al., 2007; Evi et al., 2007). This high prevalence could be due to the place and living standard of study subjects or due to a reflection of the local endemicity and geographic condition of the study area. This is almost similar to the study done around Lake Langano 83.8% (Legesse and Erko, 2004). In contrast to Seychelles, north Madagascar 94.4% (Albonico et al., 1996) the prevalence of present study was very low.

In the current study, the prevalence rate of *A. lumbricoides* (48%) was found to be higher than studies conducted in Ethiopia around lake Langano (6.2%), Dembia plains (41.3%), South Wollo (18.3%), North Gondar Dembia District (34.4%) and Adarkay (43%) (Melakebirhan et al., 1995; Jemaneh, 1997; 1998; Assefa et al., 1998; Legesse and Erko, 2004) and other countries Pakistan (4%), Cote d'Ivoire (10.8%) and Guinea (18.9%) (Glickman et al., 1999; Ahsan et al., 2005; Agbolade et al., 2007). Environmental sanitation and personal hygiene of study subjects probably play an important role for the higher prevalence rate of Ascariasis. In contrast, lower in the studies conducted in Nigeria (53.4%) and south Ethiopia Wondo Genet (84.3%) (Erko and Medhin, 2003;

Agbolade et al., 2007).

The second and third most prevalent intestinal parasites in this study were *G. lamblia* (41.9%) and *E. histolytica/dispar* (27.3%) respectively. This rate falls just above the upper range for nationwide prevalence rate of amoebiasis and giardiasis (Tesfa-Yohannes and Kloos, 1988). This could be due to the study season, poor environmental sanitation, water source of the study subjects and the favorable environmental condition of the study area for the multiplication of these parasites. However, which need further investigation.

In the present study the prevalence rate of *S. mansoni* (15.9%) was higher than compared to the previous study done in Babile 4.3%, Dembia district 5.2% (Jemaneh, 1998; Tadesse, 2005). The reason could be due to the place of the study subjects, which were found near shore of Lake Tana and the behavioral factors of the study subjects (such as frequent contact, drinking and swimming) in the lake. In contrast, much lower in the studies conducted in other areas of the country (Jemaneh, 1997; 1998; Assefa et al., 1998; Tiruneh et al., 2001; Legesse and Erko, 2004). However, it was comparable to the study done in south Gondar (14.6%) (Jemaneh, 2000).

On the other hand, the present study showed that the prevalence rate of the remaining intestinal parasite infection varies in some areas and also comparable in other study areas. The differences in prevalence rate between the study areas might be due to culture, practice, living standard and category of the study population in addition to the period of study and the methods employed for stool examination. In localities where numerous kinds of intestinal parasites found, multiple infection was frequently encountered. In this study single, double, triple and quadruple infection (13.6, 43.9, 22 and 0.3%) were respectively observed. The most frequent mixed infections in some areas were infections which involve *A. lumbricoids*, *T. trichuria* and Hookworm (Chandrasekhar and Nagesha, 2003; Agbolade et al., 2007; Evi et al., 2007). In contrast, *A. lumbricoids* and *G. lamblia* tends to appear the most frequent intestinal parasites in this study. This is probably due to environmental conditions that was favorable for the two parasites live together or the occurrence of high prevalence of the two parasites in the study area.

The present study also assessed the possible association of intestinal parasite infection with potential risk factors among school children. Several recent studies have identified a range of environmental, behavioral and social risk factors associated with intestinal parasite infections (Tesfa-Yohannes and Kloos, 1988; Erko et al., 1995; Wördemann et al., 2006).

However, very few were significantly associated in this study making it comparable to earlier studies in Ethiopia and other less developed countries. One of the factors strongly associated with intestinal parasite infection in

this study was low educational level of children mothers'. This finding is similar with the results of other study (Pinar et al., 2004; Wördemann et al., 2006). This is more likely that parents of children at high level of education provide better sanitation condition for their children than low educational level parents. Significantly higher prevalence of intestinal parasite infection also found among subjects who were not washed their hands before meal compared to those who were washed their hands regularly. This is probably due to low knowledge of children about the feco-oral transmission of intestinal parasite through their unwashed hands. The other factor that exposed children for intestinal parasite infection identified in this study was eating of unwashed /uncooked vegetables. The reason might be due to the contamination of vegetables with fecal materials in the farm. Similar findings also found in other studies growing of vegetables in faecally-polluted gardens were all found to be conducive for transmission of geohelminthes and intestinal protozoa (Erko et al., 1995).

The present study also found that using water from a river and unprotected well/spring were risk factors for intestinal parasite infection in children. This may arise from the contamination of water with animals and human waste that flooding in to the river or unprotected spring. However, other unknown factors may contribute to the increased risk associated with river/spring water and merit further investigation.

## Conclusion

Intestinal parasitic infection was highly prevalent and important health problem among Delgi School children. Therefore, an intervention strategy should be designed and implemented including provision of adequate and safe water supply, regular deworming and health education on personal hygiene to the students and to the parents.

## ACKNOWLEDGEMENTS

We would like to thank Amahra Regional Health Bureau for the provision of drugs to treat intestinal parasite infected children. We are grateful to the Delgi school community and staff of Delgi Health Center for their enthusiastic co-operation. University of Gondar and Addis continental institute of public health provided proper working conditions for the smooth running of this study also acknowledged.

## REFERENCES

Agbolade OM, Agu NC, Adesanya OO, Odejaye AO, Adigun AA, Adesanlu EB, Ogunleye FG, Sodimu AO, Adeshina SA, Bisiriyu GO,

- Omotoso OI, Udia KM. (2007). Intestinal helminthes and schistosomiasis among school children in an urban center and some rural communities in south west Nigeria. *Korean J. Parasitol.*, 45: 233-238.
- Ahsan-ul-W, Abdul B, Aziz ur R, Khawaja FQ (2005). Frequency of Intestinal Parasite infestation in children Quetta Hospital. *Pakistan J. Med. Res.*, 44: 87-88.
- Albonico M, Shamlaye N, Shamlaye C, Savioli L (1996). Control of intestinal parasitic infection in Seychelles. *Bull. World Health Organ.*, 74: 577-586.
- Assefa T, Dejene A, Michael WT (1998). Intestinal parasitism among students in three localities in south wollo Ethiopia. *Ethiop. Med. J.*, 12: 231-235.
- Chandrasekhar MR, Nagesha CN (2003). Intestinal helminthic infestation in children. *Ind. J. Pathol. Microbiol.*, 46: 492-494.
- Delgi Town Municipality Record (2009). North Gondar, Ethiopia.
- Erko B, Medhin G (2003). Human Helminthiasis in Wondo genet Southern Ethiopia with emphasis on geo helminthiasis. *Ethiop. Med. J.*, 41: 333-334.
- Erko B, Medhin G, Birrie H (1995). Intestinal parasitic infection in Bahir Dar and risk factors for transmission. *Trop. Med.*, 37: 73-78.
- Evi JB, Yavo W, Barro-kikpc, Menan EH, Koné M (2007). Intestinal helminthosis in school back ground in six towns of south western Coted ivoire. *Bull. Soc. Pathol. Exot.*, 100: 176-177.
- Glickman LT, Camara AO, Glickman NW, McCabe GP (1999). Nematode intestinal parasites of children in rural Guinea, prevalence and relationship to geophagia. *Int. J. Epidemiol.*, 28: 169-174.
- Hadidjaja P, Bonang E, Suyardi MA, Abidin SA, Ismid IS, Margono SS (1998). The effect of intervention methods on nutritional status and cognitive function of primary school children infected with *Ascaris lumbricoides*. *Am. J. Trop. Med. Hyg.*, 59: 791-795.
- Jemaneh L (1997). Intestinal helmenthic infections in school children Adarkay woreda (district) North West Ethiopia with special reference to *Schistosoma mansoni*. *Ethiop. J. Health Dev.*, 11: 289-294.
- Jemaneh L (1998). *Schistosoma mansoni* and geo-helminthiasis in school children in Dembia plains North West Ethiopia. *Ethiop. J. Health Dev.*, 12: 237-244.
- Jemaneh L (2000). The epidemiology of *Schistosoma mansoni* and soil-transmitted helminthes in elementary school children from the South Gondar Zone of the Amhara National Regional State Ethiopia. *Ethiop. Med. J.*, 38: 105-118.
- Kloos H, Tesfa-Yohannes T (1993). Intestinal parasitism in Ethiopia. In: Kloss H, Zein AZ, (editors). *The ecology of health and disease in Ethiopia*. Oxford: West view Press, pp. 223-235.
- Legesse M, Erko B (2004). Prevalence of intestinal parasite among school children in rural area close of south east of Lake Langanu. *Ethop. J. Health Dev.*, 18: 116-20.
- Magambo JK, Zeyhle E, Wachira TM (1998). Prevalence of intestinal parasite among children in south Sudan. *East Afr. Med. J.*, 75: 288-290.
- Melakebirhan D, Hailu T, Solomon T, Jilalu A, Fereju M (1995). Intensity of infection and reinfection rate of *Ascaris lumbricodes* in a rural farming village Dembia district North West Ethiopia. *Ethiop. Med. J.*, 5: 233-243.
- Merid M, Hegazy M, Mekete G, Tekle-mariam S (2001). Intestinal helminthic infection among children at lake Awassa area, south Ethiopia. *Ethiop. J. Health Dev.*, 15: 31-37.
- Nokes C, Bundy DA (1994). Does helminthes infection affect mental processing and educational achievement? *Parasitol. Today*, 10: 14-18.
- Nokes C, Grantham-McGregor SM, Sawyer AW, Cooper ES, Robinson BA, Bundy DA (1992). Moderate to heavy infection of *Trichuris trichuria* affects cognitive function in Jamaican school children. *Parasitol. Today*, 104: 539- 547.
- Phiri K, Whitty CJ, Graham SM, Ssembatya-Lule G (2000). Urban/rural distance in prevalence of intestinal helminthes in southern Malawi. *Ann. Top. Med. Parasite*, 94: 381-387.
- Pinar O, Sema E, Berna G, Ozlem O, Erdal B (2004). Intestinal parasites prevalence and related factors in school children, a western city sample-Turkey. *BMC Pub. Health*, 4: 64.
- Stephenson LS, Latham MC, Adams EJ, Kinoti SN, Pertet A (1993). Physical fitness, growth and appetite of Kenyan school boys with hookworm, *Trichuris trichiura* and *Ascaris lumbricoides* infections are improved four months after a single dose of albendazole. *J. Nutr.*, 123: 1036-1046.
- Tadesse G (2005). Prevalence of intestinal helminthic infection associated risk factors among school children in Babile town eastern Ethiopia. *Ethiop. J. Health Dev.*, 19: 140-147.
- Tesfa-Yohannes T, Kloos H (1988). Intestinal parasitism. In: Zein A.Z and Helmut Kloos (Eds). *The ecology of health & disease in Ethiopia*. Addis Ababa: Ministry of Health, pp. 214 -230.
- Tiruneh M, Fantahun M, Kassu A, Tiruneh G, Van Lieshout L (2001). *Schistosoma mansoni* in school attenders and non- attenders in North West Ethiopia. *Ethiop. J. Health Dev.*, 15: 117-123.
- Udonsi JK, Amabibi MI (1992). The human environment occupation and possible water born transmission of Hookworm in endemic coastal communities of Nigeria. *Pub. Health*. 106: 63-71.
- Wolde-michael T, Assefa T, Seyom S (1990). Intestinal parasitism among the student population of the wanji showa sugar state. *Ethiop. J. Health Dev.*, 4: 45-49.
- Wördemann M, Polman K, Menocal Heredia LT, Diaz RJ, Madurga AM, Núñez Fernández FA, Cordovi Prado RA, Espinosa AR, Duran LP, Gorbea MB, Rivero LR, Gryseels B (2006). Prevalence of intestinal parasite infections and their risk factors in Cuba children. *Trop. Med. Int. Health*, 11: 1813-1820.
- World Health Organization (1998). *Control of Tropical Diseases*. Geneva, Switzerland.
- World Health Organization expert committee (1987). *Public Health significance of intestinal parasitic infection*. WHO Bull., 65: 575-588.
- Yeneneh H (1994). Survey of intestinal parasites in Bure area Southwestern Ethiopia. *Ethiop. J. Health Dev.*, 8: 29-35.