

Full Length Research Paper

# Phytochemical Analysis of *Anastatica hierochuntica* L. and Its Implications for Traditional Medicine

Ihsanullah DAUR

Department of Arid Land Agriculture, Faculty of Meteorology, Environment and Arid Land Agriculture, King Abdulaziz University, P. O. Box 80208, Jeddah: 21589, Saudi Arabia. E-mail: [ihsanullah.daur@yahoo.com](mailto:ihsanullah.daur@yahoo.com).  
Tel: +966-2-6400000. Fax: +966-2-6952364.

Accepted 9 February, 2025

**Kaff Maryam (*ANASTATICA HIEROCHUNTICA* L.) is believed to be very useful in Arab countries for treating various health disorders. The present research highlights the chemical composition and properties of *A. HIEROCHUNTICA* L. that may be related to its beneficial effect. Eight minerals (Mg, Ca, Cr, Mn, Fe, Co, Cu, and Zn), the total phenolic content, antioxidant activity, and free radical-scavenging activity were examined in the leaf buds, stem biomass, and root biomass of Kaff Maryam. The plant was found to be a rich source of Mg, Ca, Mn, and phenolic compounds, and had potential antioxidant and free radical-scavenging activities. The findings of the study indicated that the chemical properties of the plant may rationalize its use for the treatment of menstrual cramps, asthma, depression, headache, fatigue, depression, high blood pressure, and infertility problems, and to ease childbirth, as it is used in traditional medicine. Our findings will encourage herbalists to incorporate *A. HIEROCHUNTICA* L. in the treatment of various disorders, and this herb could be proved useful in the food industry.**

**Key words:** Kaff Maryam (*Anastatica hierochuntica* L.), minerals, phenolic compounds, antioxidant activity, free radical-scavenging activity, traditional herbal medicine.

## INTRODUCTION

*Anastatica hierochuntica* L. is a small, gray winter annual herb that grows to a maximum height of 15 cm, and produces small white flowers. It is found in the arid regions of Saudi Arabia, Egypt, Jordan, Iraq, the UAE, Iran, Israel, Kuwait, and North Africa, and can survive without water for long periods. The herb curls inward under dry conditions, and then emerges from this dormant state when water is available.

*A. hierochuntica* L. is known as Kaff Maryam in Arab countries, where it is widely consumed as a tea beverage. It is powdered, mixed with honey and taken for the treatment of many conditions, in particular as a remedy for difficult childbirth and uterine hemorrhage (Khalifa and Ahmad, 1980; El-Ghazali et al., 2010). In addition, it is used to treat asthma, gastrointestinal disorders, depression, high blood pressure, indigestion, headache, cold, fever, malaria, epilepsy, fatigue, diabetes, heart disease, and infertility (Batanouny, 1999; Eman et al., 2011). The idea for the present research

was generated from the studies of Baliga et al. (2011), Ducat et al. (2011), Vadivel et al. (2011), and Kovacik et al. (2012), who have identified and quantified a number of minerals and phenolic compounds from various plants used for treating health disorders. However, studies available on the chemical properties of Kaff Maryam are limited (El-Ghazali et al., 2010; Nakashima et al., 2010). The present study provides the first comprehensive analysis of the minerals and total phenolic contents, as well as antioxidant and free radical-scavenging activities found in various parts of the *A. hierochuntica* L. plant, and examines the herb's relation to folk medicine use.

## MATERIALS AND METHODS

### Samples preparation

*A. hierochuntica* L. plants were purchased from Makkah, Saudi Arabia. After that the plant were separated into leaf buds, stem

**Table 1.** Instrument settings for flame atomization.

Condition	Mg	Ca	Cr	Mn	Fe	Co	Cu	Zn
Wavelength (nm)	285.2	422.7	359.3	279.6	248.3	240.7	324.8	213.9
Lamp current (mA)	9.0	10	9.0	9.0	15.0	15.0	9.0	6.5
Slit width (nm)	0.2	0.2	1.3	0.4	0.2	0.2	1.3	1.3
Oxidant	Air							
Oxidant pressure (kPa)	160	160	160	160	160	160	160	160
Fuel	Air C <sub>2</sub> H <sub>2</sub>							
Fuel pressure (l/min)	2.2	2.2	2.8	2.2	2.0	2.2	2.2	2.0
Calculation mode	Integral							
Calculation time (s)	5.0	5.0	5.0	5.0	5.0	5.0	5.0	5.0

biomass, and root biomass. All the plant parts were oven-dried, and then, powdered for uniformity, and stored in an airtight sample jars at room temperature. For subsequent mineral analysis, 50 g of the powder of each individual plant part was converted to ash at 550°C in a muffle furnace, according to Daur et al. (2011).

For determining total phenolic content, antioxidant activity, and free radical-scavenging activity, 50 g of the powder of each individual plant part was extracted in triplicate with distilled water by boiling and stirring for 10 min. Then, the extracts were filtered over Whatman No. 1 filter paper and stored at -80°C until use, as described by Sharma and Gujral (2010).

#### Minerals analysis

Mineral analysis was performed according to the AOAC method (2010), where 1.0 g of the ash prepared as mentioned earlier was digested with 10.0 ml of concentrated hydrochloric acid in a beaker on a hot plate until the solution was clear; the volume was reduced to 1.0 ml. The solution was filtered through Whatman No. 1 filter paper and then diluted to 10.0 ml with distilled water. A control solution was prepared by heating hydrochloric acid without addition of the sample.

A Hitachi Z-5000 Atomic Absorption Spectrophotometer equipped with a Zeeman background corrector and data processor was used for mineral analysis of the ash and control solutions. Eight minerals (Mg, Ca, Cr, Mn, Fe, Co, Cu, and Zn) were quantified. All functions were set strictly, according to the manufacturer's instructions, by using the flame atomization technique (Table 1). All experimental procedures, ashing, digestion of ash, and measurement of minerals were performed in triplicate, at the minimum.

#### Total phenolic content

The total phenolic content was determined according to the Folin-Ciocalteu method, as described by Sharma and Gujral (2010). The results were expressed as both micrograms of gallic acid equivalents (GAE) per gram of sample and micrograms of ferulic acid equivalents (FAE) per gram of sample.

#### Antioxidant activity

The antioxidant activity was measured according to the method described by Benzie and Strain (1999), by using the ferric-reducing ability of plasma (FRAP) assay that is based on the reduction of the ferric-tripyridyltriazine (TPTZ-Fe<sup>3+</sup>) complex to its TPTZ-Fe<sup>2+</sup> form in the presence of antioxidants. The reducing power of the extracts

was evaluated according to the method described by Yen and Chen (1995).

#### Free radical-scavenging activity

The free radical-scavenging activity of the extracts for the various plant parts were evaluated using the method described by Aquino et al. (2001), which is based on the ability of the extract to scavenge the stable 2,2-diphenyl-1-picrylhydrazyl (DPPH), resulting in the bleaching of the purple color exhibited by the stable DPPH radical that is measured at an absorbance of 517 nm. Results were expressed as the percentage of inhibition of the DPPH radical.

## RESULTS AND DISCUSSION

The results of mineral analysis of the leaf buds, stem biomass, and root biomass of *A. hierochuntica* L. are presented in Table 2. The data shows that the concentrations of Mg, Ca, and Mn in *A. hierochuntica* L. were greater than their concentrations in other medicinal plants reported by Karadaş and Kara (2012). Furthermore, the concentrations of other minerals (Fe, Cu, and Zn) in *A. hierochuntica* L. were also either greater than or comparable to the concentrations in other medicinal plants, as reported by Lokhande et al. (2010) for *Vetiveria zizanioides*, Shirin et al. (2010) for *Withania somnifera*, and Karadaş and Kara (2012) for cinnamon. The minerals Fe, Cu, and Zn have important functions in blood, tissue, and plasma, and are often consumed as supplements, as reported by Agarwal et al. (2011). The medicinal value of *A. hierochuntica* L. may possibly be related to the high concentrations of Mg, Ca, and Mn contained in the plant. Magnesium has a curative effect on more than 300 disorders, including menstrual cramps, asthma, headache, fatigue, depression, diabetes, fibromyalgia, arrhythmia and heart failure, and preeclampsia and eclampsia, as well as in easing childbirth, as reported by Chiladakis et al. (2001), Champagne (2008), Sun-Edelstein and Mauskop (2009), Guerrero et al. (2009), Guerrero-Romero and Rodríguez-Morán (2009), and Chiueve et al. (2011). Calcium is required for bone repair, vascular contraction and vasodilatation, muscle function, nerve transmission,

**Table 2.** Mineral concentrations in leaf buds and entire biomass of *A. hierochuntica* L.

Test sample	Minerals (ppm)							
	Mg	Ca	Cr	Mn	Fe	Co	Cu	Zn
LEAF BUDS	8114 ± 4.6 <sup>A</sup>	23286 ± 15.2 <sup>B</sup>	5.10 ± 0.12 <sup>A</sup>	676.5 ± 11.4 <sup>A</sup>	2580.2 ± 9.4 <sup>A</sup>	1.5 ± 0.08 <sup>A</sup>	290.6 ± 6.32 <sup>A</sup>	37111.6 ± 5.2 <sup>A</sup>
STEM BIOMASS	6998 ± 9.5 <sup>B</sup>	22954 ± 21.4 <sup>B</sup>	4.38 ± 0.10 <sup>B</sup>	415.4 ± 8.6 <sup>C</sup>	2186.3 ± 5.8 <sup>B</sup>	1.4 ± 0.04 <sup>B</sup>	143.0 ± 0.03 <sup>B</sup>	23241.0 ± 5.5 <sup>B</sup>
ROOT BIOMASS	6475 ± 12.2 <sup>B</sup>	28322 ± 44.0 <sup>A</sup>	4.51 ± 0.03 <sup>B</sup>	597.1 ± 14.2 <sup>B</sup>	2138 ± 14.5 <sup>B</sup>	1.3 ± 0.10 <sup>B</sup>	156.3 ± 0.03 <sup>B</sup>	24867.0 ± 8.4 <sup>B</sup>

Each value in the table is the mean of three replicates ± standard deviation. Mean values with different superscript letters in the same row differ significantly ( $p < 0.05$ ).

**Table 3.** Total phenolic content, Antioxidant activity and free radical scavenging activity of Kaff Maryam (*A. hierochuntica* L. ).

Test sample	Total phenolic content		Antioxidant activity	Free radical scavenging activity
	GAE <sup>a</sup> (µg/g)	FAE <sup>b</sup> (µg/g)	FRAP (µM FeSO <sub>4</sub> /g)	[DPPH inhibition (%)]
Leaf buds	2193.1 ± 8.0 <sup>a</sup>	1089.4 ± 12.0 <sup>a</sup>	2864.8 ± 36.4 <sup>a</sup>	65 <sup>a</sup>
Stem biomass	1659.7 ± 16.7 <sup>b</sup>	661.0 ± 19.4 <sup>b</sup>	1738.5 ± 51.3 <sup>c</sup>	48 <sup>c</sup>
Root biomass	1680.5 ± 21.5 <sup>b</sup>	662.6 ± 10.3 <sup>b</sup>	1987.4 ± 48.5 <sup>b</sup>	58 <sup>b</sup>

Each value in the table is the mean of three replicates ± standard deviation, Mean values with different superscript letters in the same row differ significantly ( $p < 0.05$ ), <sup>a</sup> GAE, gallic acid equivalents, <sup>b</sup> FAE, ferulic acid equivalents.

intracellular signaling, and hormonal secretion, as well as for many other metabolic functions, as mentioned by Moyad (2003), Straub (2007), and Lappe et al. (2008). Moreover, Mn plays a role in the formation of connective tissue, bones, blood clotting factors, and sex hormones, and protects against certain types of cancer, as supported by Ansari et al. (2004), Son et al. (2007), and Menezes-Filho et al. (2011).

In addition to being a rich source of important minerals, all parts of *A. hierochuntica* L., especially the leaf buds, were found to be rich in total phenolic content, and to have high antioxidant and free radical-scavenging activities (Table 3). Recently, the phenolic content and the antioxidant and free radical-scavenging properties of natural products have generated tremendous research interest owing to their related anticancer and antimutagenic properties, as well as their protective effects against cardiovascular diseases and their ability to confer decreased risk of certain chronic diseases, as reported by Weichselbaum and Buttriss (2010) and Saura-Calixto (2011).

In conclusion, all parts of the *A. hierochuntica* L. plant were found to be rich in essential minerals and total phenolic, and to have high antioxidant and free radical-scavenging activities. These properties help to explain the therapeutic effect of the plant as an herbal medicine.

## ACKNOWLEDGMENTS

This paper was funded by the Deanship of Scientific Research (DSR), King Abdulaziz University, Jeddah, under grant No. (19-130-D1432). The author, therefore, acknowledge with thanks DSR technical and financial

support.

## REFERENCES

- Agarwal A, Khanna P, Baidya DK, Arora MK (2011). Trace elements in critical illness. *J. Endocrinol. Metabol.*, 1: 57-63.
- Ansari TM, Ikram N, Najam-ul-Haq M, Fayyaz I, Fayyaz Q, Ghafoor I, Khalid N (2004). Essential Trace Metal (Zinc, Manganese, Copper and Iron) Levels in Plants of Medicinal Importance. *J. Biol. Sci.*, 4: 95-99.
- AOAC (2010). Latimer G.W. (Ed.) Official Methods of Analysis of AOAC International, 18th Edition, Revision 3. AOAC International, Gaithersburg, Maryland, USA.
- Aquino R, Morelli S, Lauro MR, Abdo S, Saija A, Tomaino A (2001). Phenolic constituents and antioxidant activity of an extract of *Anthurium versicolor* leaves. *J. Nat. Prod.*, 64: 1019-1023.
- Baliga MS, Bhat HP, Baliga BRV, Wilson R, Palatty PL (2011). Phytochemistry, traditional uses and pharmacology of *Eugenia jambolana* Lam. (black plum): A review. *Food Res. Int.*, 44: 1776-1789.
- Batanouny KH (1999). *Wild Medicinal Plants in Egypt*. The Palm Press. Cairo, p. 207.
- Benzie IFF, Strain JJ (1999). Ferric reducing antioxidant power assay: direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. *Meth. Enzymol.*, 299: 379-389.
- Champagne CM (2008). Magnesium in hypertension, cardiovascular disease, metabolic syndrome, and other conditions (A review). *Nutr. Clin. Pract.*, 23: 142-151.
- Chiladakis JA, Stathopoulos C, Davluros P, Manolis AS (2001). Intravenous magnesium sulfate versus diltiazem in paroxysmal atrial fibrillation. *Int. J. Cardiol.*, 79: 287-291.
- Chiuve SE, Korngold EC, Januzzi JL, Gantzer ML, Albert CM (2011). Plasma and dietary magnesium and risk of sudden cardiac death in women. *Am. J. Clin. Nutr.*, 93: 253-260.
- Daur I, Sepetoğlu H, Sindel B (2011). Dynamics of faba bean growth and nutrient uptake and their correlation with grain yield. *J. Plant Nutr.*, 34: 1360-1371.
- Ducat G, Torres YR, Dalla-Santa HS, Caetano IK, Kleinubing SA,

- Stock D, Tussolini L, Justo TH, Quinaia SP (2011). Correlation among metallic ions, phenolic compounds and antimicrobial action in medicinal plants extracts. *J. Food Qual.*, 34: 306-314.
- El-Ghazali GE, Al-Khalifa KS, Saleem GA, Abdallah EM (2010). Traditional medicinal plants indigenous to Al-Rass province, Saudi Arabia. *J. Med. Plant Res.*, 4: 2680-2683.
- Eman AS, Tailang M, Benyounes S, Gauthaman K (2011). Antimalarial and hepatoprotective effects of entire plants of *Anastatic hierochuntica*. *Int. J. Res. Phytoch. Pharm.*, 1: 24-27.
- Guerrera MP, Volpe SL, Mao JJ (2009). Therapeutic uses of magnesium. *Am. Fam. Phys.*, 80: 157-162.
- Guerrero-Romero F, Rodríguez-Morán M (2009). The effect of lowering blood pressure by magnesium supplementation in diabetic hypertensive adults with low serum magnesium levels: a randomized, double-blind, placebo-controlled clinical trial. *J. Hum. Hypertens.*, 23: 245-251.
- Karadağ C, Kara D (2012). Chemometric approach to evaluate trace metal concentrations in some spices and herbs. *Food Chem.*, 130: 196-202.
- Khalifa TI, Ahmad MA (1980). Pharmacognostical study of certain species of *Anastatica*. PhD thesis, University of Cairo, Egypt.
- Kovacik J, Gruz J, Klejdus B, Stork F, Hedbavny J (2012). Accumulation of metals and selected nutritional parameters in the field-grown chamomile *anthodia*. *Food Chem.*, 131: 55-62.
- Lappe J, Cullen D, Haynatzki G, Recker R, Ahlf R, Thompson K (2008). Calcium and vitamin D supplementation decreases incidence of stress fractures in female navy recruits. *J. Bone Min. Res.*, 23: 741-749.
- Lokhande R, Singare P, Andhale M (2010). Study on Mineral content of Some Ayurvedic Indian Medicinal Plants by Instrumental Neutron Activation Analysis and AAS Techniques. *Health Sci. J.*, 4: 157-168.
- Menezes-Filho JA, Novaes CO, Moreira JC, Sarcinelli PN, Mergler D (2011). Elevated manganese and cognitive performance in school-aged children and their mothers. *Environ. Res.*, 111: 156-163.
- Moyad MA (2003). The potential benefits of dietary and/or supplemental calcium and vitamin D. *Urol. Oncol.*, 21: 384-391.
- Nakashima S, Matsuda H, Oda Y, Nakamura S, Xu F, Yoshikawa M (2010). Melanogenesis inhibitors from the desert plant *Anastatica hierochuntica* in B16 melanoma cells. *Bioorg. Med. Chem.*, 18: 2337-2345.
- Saura-Calixto F (2011). Dietary fibre as a carrier of dietary antioxidants: an essential physiological function. *J. Agric. Food Chem.*, 59: 43-49.
- Sharma P, Gujral HS (2010) Antioxidant and polyphenol oxidase activity of germinated barley and its milling fractions. *Food Chem.*, 120: 673-678.
- Shirin K, Imad S, Shafiq S, Fatima K (2010). Determination of major and trace elements in the indigenous medicinal plant *Withania somnifera* and their possible correlation with therapeutic activity. *J. Saudi Chem. Soc.*, 14: 97-100.
- Son EW, Lee SR, Choi HS, Koo HJ (2007). Effects of supplementation with higher levels of manganese and magnesium on immune function. *Arch. Pharm. Res.*, 30: 743-749.
- Straub DA (2007). Calcium supplementation in clinical practice: a review of forms, doses, and indications. *Nutr. Clin. Prac.*, 22: 286-296.
- Sun-Edelstein C, Mauskop A (2009). Foods and supplements in the management of migraine headaches. *Clin. J. Pain*, 25: 446-452.
- Vadivel V, Nandety A, Biesalski HK (2011). Antioxidant, free radical scavenging and type II diabetes-related enzyme inhibition properties of traditionally processed Jequirity bean (*Abrus precatorius* L.). *Int. J. Food Sci. Tech.*, 46: 2505-2512.
- Weichselbaum E, Buttriss JL (2010). Polyphenols in the diet. *Nutr. Bull.*, 35: 157-164.
- Yen GC, Chen HY (1995). Antioxidant activity of various tea extracts in relation to their antimutagenicity. *J. Agric. Food Chem.*, 43: 27-32.