

Full Length Research Paper

Salt Stress Mitigation in Chickpea by Exogenous Application of Ascorbic Acid

Mohamed Salah Beltagi

Botany Department, Faculty of Science, Suez Canal University, Ismailia, Egypt. E-mail: msbeltagi@yahoo.com.

Accepted 10 April, 2025

The effect of exogenous supply of ascorbic acid on growth and metabolic changes in chick pea under salinity treatment was carried out in pot experiment. The exogenous supply of ascorbic acid (4 mM) improved the fresh and dry matter gain in roots, shoots and leaves of chick pea (*Cicer arietinum* L.) Plants attenuated by the salt stress (40 mM NaCl) environment in the soil solution. Significant synergistic effect between NaCl (40 mM) and ascorbic acid treatment increased the contents of chlorophyll a and chlorophyll stability index (CSI %) in leaves of chick pea plants. The total number of protein bands/lane did not change under the low (20 mM) NaCl concentration but was dramatically reduced by the high (40 mM) NaCl treatment. The sum of optical densities (O. D.) of protein bands was inhibited by both levels of NaCl, but was induced by 10.68% by the added ascorbic acid at 20 mM NaCl and by 21.39% at 40 mM NaCl. Six different polypeptides of molecular weights 146.28, 117.98, 51.55, 49.6, 44.49 and 38.34 were completely disappeared under NaCl stress (40 mM). These bands reappeared in response to the added ascorbic acid treatment. Moreover, the optical density of every individual protein band was induced by ascorbic acid under the low NaCl concentration. The results indicate synergistic interaction between salinity stress and ascorbic acid for the sake of salt resistance in chick pea plants.

Keywords: Plant growth, photosynthetic pigments, chlorophyll stability index, SDS-PAGE protein.

INTRODUCTION

Environmental stresses trigger a wide variety of plant responses, ranging from altered gene expression and cellular metabolism to accelerated leaf senescence and permanent wilting, all lead to changes in growth rates and crop yield. Salt stress adversely affects plant growth and productivity (Reddy et al., 2004; Baek et al., 2005; Amor et al., 2005), through the increase in reactive oxygen species (ROS) which may cause oxidative stress resulting in cellular damage by oxidation of lipids, proteins and nucleic acids (McKersie and Leshem, 1994; Pastori and Foyer, 2002; Apel and Hirt, 2004). To minimize the effects of oxidative salt stress, plant cells have evolved a complex antioxidant system, which is composed of low-molecular mass antioxidants as well as ROS-scavenging enzymes (Alscher et al., 1997; Apel and Hirt, 2004).

One approach for inducing oxidative stress tolerance would be to increase the cellular level of enzyme substrates such as ascorbic acid (vitamin C). Ascorbic acid is a small, water-soluble anti-oxidant molecule which

acts as a primary substrate in the cyclic pathway of enzymatic detoxification of hydrogen peroxide. Improved understanding of ascorbate in plants will lead to the possibility of increasing ascorbate concentration in plants by genetic manipulation. This will have benefits for tolerance of plants to oxidative stresses (Foyer, 1993; Smirnoff, 1995).

Salinity is generally detrimental to plant growth through its adverse effects on plant metabolism that induces important modifications in gene expression. Such modifications may lead to accumulation or depletion of certain metabolites resulting in an imbalance in the levels of a relatively small set of cellular proteins, which could increase, decrease, appear or disappear after salt treatment (Kong-ngern et al, 2005).

This study investigates the effects of the exogenous supply of ascorbic acid on the growth and anabolic changes of chick pea plants with a special emphasis on the role of the cellular levels of SDS-PAGE protein profile in relation to salt tolerance.

Table 1. Growth parameters of chick pea (*Cicer arietinum* L.) plants treated with 20 mM NaCl (T1), 40 mM NaCl (T3), 20 mM NaCl + 4 mM ascorbic acid (T2) and 40 mM NaCl + 4 mM ascorbic acid (T4), respectively. C, controlled plants.

Treatment	Stem length (cm)	Root length (cm)	No. of leaves/ plant	Stem f. wt. (g)	Root f. wt. (g)	Stem d. wt. (g)	Root d. wt. (g)
C	24.33	12.50	19.67	28.67	6.72	7.33	0.433
T ₁	23.67	11.00	15.33	27.91	5.67	6.63	0.336
T ₂	28.67	16.33	19.00	30.19	6.44	8.33	0.322
T ₃	21.75	6.67	14.67	25.80	4.33	6.11	0.288
T ₄	23.33	6.83	17.33	29.67	5.11	7.21	0.367
LSD ($P \geq 0.05$)	5.950	7.515	4.052	3.055	1.801	1.476	0.101

MATERIALS AND METHODS

Plant material and treatments: Pure strain of chick pea (*Cicer arietinum* L.) seeds were surface-sterilized with 15% (w/v) sodium hypochlorite for 20 min, rinsed with distilled water, then soaked in plastic pots (15 cm) in pre-sieved homogenous garden soil (sandy loam). All planted pots were kept in the open garden in about 31/22°C day/night temperature and average relative humidity 60% and irrigated regularly up to field capacity with pure water. After two weeks, planted pots were randomly subdivided into five equal groups (5 pots each). One group were kept irrigated with pure water and sampled as control. Groups 2 and 3 were subjected to 20 and 40 mM salinity (NaCl); while groups 4 and 5 received the same salinity treatment plus 4 mM ascorbic acid, respectively. Upon treatments, all pots received equal quantities of 1/2 Hoagland nutrient solution.

Growth parameters: By the end of the experiment (8 weeks), plants were smoothly uprooted, cleaned from soil residues and prepared for measurements. Plant height, root length, shoot and root fresh and dry weights, total number of leaves and number of lateral branches per plant were recorded.

Estimation of photosynthetic pigments: Chlorophyll a, chlorophyll b and carotenoids ($\mu\text{g/ml}$) contents was estimated in the fresh leaves in 80% acetone according to the procedure described by Lichtenthaler and Wellburn (1983). Chlorophyll stability index (CSI %) was calculated by combining chlorophyll a+b contents in chick pea foliage leaves before and after salinity stress following the formula noted by Kumari et al. (2004): $\text{CSI \%} = \frac{\text{Chlorophyll before stress} - \text{chlorophyll under stress}}{\text{chlorophyll under stress}} \times 100$.

Protein extractions: For SDS-PAGE protein extracts were prepared by extracting appropriate weight from the frozen plant material with 0.125 M tris/ borate, pH 8.9. All the obtained extracts were kept at 4°C for 24 h and centrifuged at 10000 rpm for 20 min. The supernatants were used for electrophoresis.

SDS-PAGE protein analysis: Exact 5 μg of each sample was separated on gel slabs according to the method of Laemmli (1970). Protein subunit bands were stained with Coomassie blue R- 250 by standard techniques. The gel was scanned using Gel pro- Analyzer ver. 3.3 (Media Cybernetics 93-97).

Statistical analysis: Data were statistically analyzed by multiple comparison procedure at $p \leq 0.05$ using t-test and mean separation by least significant difference (LSD) (Steel and Torrie, 1980).

RESULTS

Plant growth: The statistical analyses of growth parameters (Table 1) revealed non significant effect on stem length, root length and stem dry weight at all salinity or salinity + ascorbic acid treatment. However, the stem and root fresh weights and the root dry weight were inhibited by the higher salinity treatment (40 mM NaCl). The number of leaves per plant was inhibited by both salinity levels (20 mM and 40 mM NaCl). The addition of ascorbic acid (4 mM) significantly increased the stem dry weight at the low level of NaCl (20 mM); while the stem fresh weight and the root dry weight were increased at the higher level of NaCl (40 mM).

Photosynthetic pigments: The spectrophotometric estimation of chlorophyll pigments (Table 2) indicated significant inhibition of chlorophyll a content (mg/g) at the high NaCl concentration (40 mM). This inhibition was recovered by the added ascorbic acid (4 mM). The content of chlorophyll b did not show significant changes under any of the treatments. The total chlorophylls a + b was not inhibited by the applied salinity levels, but increased in response to the added ascorbic acid. On the other hand, the value of chlorophyll stability index (CSI%) was higher at the combination of 40 mM NaCl + 4 mM ascorbic acid (Figure 1).

Protein analysis: The SDS-PAGE analysis of the protein profile of chick pea plants revealed both qualitative (Figures 2 and 3) and quantitative (Table 3) changes in the banding patterns of proteins. The total number of protein bands/lane did not change under the low (20 mM) NaCl concentration but was dramatically reduced by the high (40 mM) NaCl treatment. The sum of optical densities (O. D.) was inhibited by both levels of NaCl. Six different polypeptides of molecular weights 146.28, 117.98, 51.55, 49.6, 44.49 and 38.34 were completely disappeared in response to NaCl (40 mM) stress. These bands reappeared in response to the added ascorbic acid treatment (4 mM). Moreover, the optical density of every individual protein band was induced by ascorbic acid (4 mM) under the low NaCl concentration (20 mM). The sum

Table 2. Chlorophyll pigments ($\mu\text{g/ml}$) in chick (*Cicer arietinum* L.) leaves treated with 20 mM NaCl (T1), 40 mM NaCl (T3), 20 mM NaCl + 4 mM ascorbic acid (T2) and 40 mM NaCl + 4 mM ascorbic acid (T4), respectively. C, controlled plants.

Treatment	Chl a	Chl b	Chl a + Chl b	Chl a/Chl b
C	0.295	0.112	0.408	2.63
T ₁	0.232	0.077	0.308	3.02
T ₂	0.274	0.120	0.394	2.27
T ₃	0.211	0.092	0.312	2.32
T ₄	0.325	0.158	0.484	2.05
LSD ($P \geq 0.05$)	0.081	0.054	0.129	0.659




Figure 1. Chlorophyll Stability Index (CSI %) in chick pea (*Cicer arietinum* L.) plants subjected to: 1) 20 mM NaCl + 4 mM ascorbic acid and 2) 40 mM NaCl + 4 mM ascorbic acid.




Figure 2. Electrophotograph SDS-PAGE of total proteins of chick pea (*Cicer arietinum* L.) plants. Track 1, untreated (controlled) plants; tracks 2 and 4, plants treated with 20 and 40 mM NaCl; tracks 3 and 5, plants treated with 20 mM NaCl + 4 mM ascorbic acid and 40 mM NaCl + 4 mM ascorbic acid, respectively. Track M, molecular weight markers used on polyacrylamide gel.

of O. D. was induced by 10.68% by the added ascorbic acid at the low level of salt stress and by 21.39% at the high salt concentration indicating a synergistic influence of salinity stress and ascorbic acid on salt-stressed chick pea plants.

DISCUSSION

In the present investigation, the responses of chick pea plants to high levels of salinity were reflected by decreases in stem fresh weight, root fresh weights, root dry weight and the number of leaves per plant. The stressful environment in the soil solution at a concentration of 40 mM NaCl attenuated the fresh and dry matter gain in roots, shoots and leaves. The inhibitory effects of salt stress on these parameters add more support to the ubiquitous findings of earlier investigations (Pérez- Alfocsa et al., 1993; Hamada, 1996). The reduced plant growth under salt stress conditions could be attributed to the physiological drought induced by the low water potential of the soil solution and osmotic adjustments in plants as a result of increased ionic concentration in their

Table 3. Comparative analysis of optical density (O. D.), molecular weight (M.Wt.) and relative front (R_f) of SDS-PAE protein profile of chick pea (*Cicer arietinum* L.) plants treated with NaCl or NaCl + ascorbic acid.

Band number	Treatment & O. D.					R _f	Mol. Wt. (KDa)
	1	2	3	4	5		
1	65.21	49.87	80.39	48.54	57.39	0.17	521.92
2	76.01	54.80	99.86	-	67.34	0.29	146.28
3	75.43	50.11	97.14	-	55.38	0.31	117.98
4	80.81	54.51	101.01	68.75	57.56	0.38	110.25
5	90.01	74.42	108.24	74.25	69.25	0.43	104.85
6	114.62	84.83	120.90	200.05	78.21	0.55	086.37
7	227.20	197.88	231.56	157.85	208.31	0.66	066.36
8	209.61	125.07	196.29	130.35	125.92	0.73	57.00
9	156.89	102.47	-	-	99.17	0.78	51.55
10	-	-	-	-	105.44	0.80	49.60
11	141.59	137.43	-	147.37	108.83	0.85	46.40
12	186.00	207.26	207.26	-	170.24	0.88	44.49
13	-	-	-	178.55	173.23	0.93	41.13
14	233.05	189.79	249.21	-	139.56	0.97	38.34
Sum of O. D.	1656.43	1328.44	1491.86	1005.71	1220.04		
Bands/lane	12	12	10	8	14		

1 = untreated (control) plants

2 and 4 = plants treated with 20 mM NaCl and 40 mM NaCl; respectively

3 and 5 = plants treated with 20 mM NaCl + 4 mM ascorbic acid and 40 mM NaCl

+ 4 mM ascorbic acid; respectively.

their cells, which result in deformation of macromolecules by disrupting their shell or bound water (Schwarz, 1985).

The biochemical functions of ascorbate have been divided into four categories, (1) antioxidant, regenerates the lipophilic antioxidant α -tocopherol, vitamin E (Asada, 1994); (2) enzyme cofactor for hydroxylase enzymes involved in the synthesis of hydroxyproline-rich glycol-proteins, cell wall structural proteins (Carpita and Gibeau, 1993); (3) electron transport, acts as an in vitro electron donor and acceptor in transmembrane electron transport (Asard et al., 1995); and (4) oxalate and tartaric acid synthesis (Saito, 1996). The results of the current research indicated that the added ascorbic acid (4 mM) improved the stem and root fresh and dry weights of salt-stressed chick pea plants. Consistent findings reported on the beneficial effects of the exogenous application of ascorbic acid in mitigating partially the adverse effects of salt stress on growth, like cell division and cell enlargement (Mozafar and Oertli, 1992; Ahmed-Hamad and Monsaly, 1998). Moreover, in tomato seedlings, the exogenous supply of ascorbic acid increased tissue levels (Arrigoni et al., 1997) and percentage age for surviving the toxic effects of NaCl (Shalata and Neumann, 2001). Alternative organic carbon sources without direct antioxidant activity did not provide equivalent protection. These remarkable protective effects appeared to be specifically related to the anti-oxidant activity of ascorbic acid, rather than its possible utility as an organic substrate for respiratory energy metabolism (Shalata and

Neumann, 2001).

Ascorbate in plants occurs in the cytosol, chloroplasts, vacuoles, mitochondria and cell wall (Anderson et al., 1983; Rautenkrantz et al., 1994). The concentration in chloroplast can be high (up to 50 mM in spinach) and is probably related to its central role in photosynthesis (Foyer, 1993). Our results revealed significant synergistic effect between NaCl (40 mM) stress and exogenous ascorbic acid (4 mM) on the contents of chlorophyll a and chlorophyll stability index (CSI %) in the leaves of chick pea plants. Since salt stress can lead to oxidative stress through the increase in reactive oxygen species (ROS) which are highly reactive and may cause cellular damage, one of the proposed biochemical modes of ascorbate is to act as an antioxidant by scavenging hydrogen peroxide (chloroplasts lack catalase) as it forms (Miyake and Asada, 1992).

A finding of the current research is the role of the exogenous ascorbic acid in salt-stressed chick pea plants and its ability in the induced stability of protein synthesis. The values of the band intensities (O.D.) were always higher in the low salt-stressed plants treated with ascorbic acid than in the only salt-stressed plants. In addition, the inhibited protein bands in the high salt-stressed individuals were completely restored by the added ascorbic acid. This enhanced protein stability could be partially attributed to the increased activity of antioxidant enzymes against salinity stress environment (Inzé and Van Montague, 1995; Noctor and Foyer, 1998;




Figure 3. Scan of the tracks in the electrophotograph (Figure 2) of SDS-PAGE of total proteins of chick pea (*Cicer arietinum* L.) plants. 1, molecular weight markers used on polyacrylamide gel; 2, untreated (controlled) plants; 3 and 5, plants treated with 20 and 40 mM NaCl; 4 and 6, plants treated with 20 mM NaCl + 4 mM ascorbic acid and 40 mM NaCl + 4 mM ascorbic acid, respectively.

McKersie et al., 1999).

In conclusion, this study shows, for the first time, the remarkable synergistic interaction between the exogenous supply of ascorbic acid and the anabolic capacity of

protein in salt-stressed chick pea plants. Molecular genetics studies should complement the biochemical and physiological approaches concerning the role of ascorbate in protein expression for increased plant resistance

to salt stress.

REFERENCES

- Ahmed-Hamad AM, Monsaly HM (1998). Seed soaking presowing in vitamins versus the adverse effects of NaCl salinity on photosynthesis and some related activities of maize and sunflower plants. In: Proceedings of the XIth International photosynthesis Conference, Budapest, Hungary, August, pp. 17-22.
- Alscher RG, Dnahue JL, Cramer CL (1997). Reactive oxygen species and antioxidants: relationship in green cells. *Physiol. Plant.* 100: 224-233.
- Amor NB, Hamed KB, Ahmed D, Grigon C, Chedly-Abdelly (2005). Physiological and antioxidant responses of the potential halophyte *Crithmum maritimum* to salinity. *Plant Sci.* 168(4): 889-899.
- Anderson JW, Foyer CH, Walker DA (1983). Light-dependent reduction of dehydroascorbate and uptake of exogenous ascorbate by spinach chloroplast. *Planta* 158: 442-450.
- Apel K, Hirt H (2004). Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Ann. Rev. Plant Biol.* 55: 373-399.
- Arrigoni O, Calabrese G, De Gara L, Bitonti MB, Liso R (1997). Correlation between changes in ascorbate and growth of *Lupinus albus* seedlings. *J. Plant Physiol.* 150: 302-338.
- Asada K (1994). Mechanisms for scavenging reactive molecules generated in chloroplasts under light stress. In: Baker NR, Bowyer JR eds. *Photoinhibition of photosynthesis. From molecular mechanism to the field.* Oxford: Bios Scientific Publishers, 129-142.
- Asard H, Horemans N, Caubergs RJ (1995). Involvement of ascorbic acid and a b-type cytochrome in plant plasma membrane redox reactions. *Protoplasma* 184: 36-41.
- Baek MLI, Kim JII, Chung BY, Kim JS, Lee IS (2005). Alleviation of salt stress by low dose gamma-irradiation in rice. *Biol. Plant.* 49(2): 273-276.
- Carpita NC, Gibeaut MD (1993). Structural models of primary cell walls in flowering plants: consistency of molecular structure with the physical properties of the walls during growth. *Plant J.* 3: 1-30.
- Foyer CH (1993). Ascorbic acid. In: Alscher RG, Hess JL eds. *Antioxidants in higher plants.* Boca Raton: CRC Press 31-58.
- Hamada AM (1996). Effect of NaCl, water stress or both on gas exchange and growth of wheat. *Biol. Plant.* 38: 405-412.
- Inzé D, Van Montague M (1995). Oxidative stress in plants. *Current Opinion Biotech.* 6: 153-158.
- Kong-ngern K, Daduang S, Wongkham C, Bunnag S, Kosittakun M, Theerakulpisut P (2005). Protein profile in response to salt stress in leaf sheaths of rice seedlings. *Sci. Asia* 31: 403-408.
- Kumari M, Sam MD, Virnala Y, Pawan A (2004). Physiological parameters governing drought in maize. *Indian J Plant Physiol.* 9(2): 203-207.
- Laemmli UK (1970). Cleavage of structural proteins during the assembly of head of bacteriophage T4. *Nature* 227: 680-685.
- Lichtenthaler HK, Wellburn RR (1983). Determination of total carotenoids and chlorophylls a and b of leaf extracts in different solvents. *Biochem. Soc. Transact.* 11: 591-592.
- McKersie BD, Booley SR, Jones KS (1999). Winter survival of transgenic alfalfa overexpressing superoxide dismutase. *Plant Physiol.* 119: 839-847.
- McKersie BD, Leshem YY (1994). *Stress and stress coping in cultivated plants.* Kluwer Academic Publishers, London.
- Miyake C, Asada K (1992). Thylakoid bound ascorbate peroxidase in spinach chloroplasts and photoreduction of its primary oxidation product, monodehydroascorbate radicals in the thylakoids. *Plant Cell Physiol.* 33: 541-553.
- Mozafar A, Oertli JJ (1992). Uptake and transport of thiamin (vitamin B1) by barley and soybean. *J. Plant Physiol.* 436: 442.
- Noctor G, Foyer CH (1998). Ascorbate and glutathione: keeping active oxygen under control. *Ann. Rev. Plant Physiol. & Molec. Biol.* 49: 249-279.
- Pastori GM, Foyer CH (2002). Common components, networks, and pathways of cross-tolerance to stress. The central role of 'redox' and abscisic acid-mediated controls. *Plant Physiol.* 129: 7460-7468.
- Pérez-Alfocea F, Estan MT, Caro M, and Guerrier G (1993). Osmotic adjustment in *Lycopersicon esculentum* and *L. pennellii* under NaCl and polyethylene glycol 6000 iso-osmotic stress stresses. *Physiol. Plant.* 87: 493-498.
- Rauten-kranz AAF, Li L, Machler F, Martinoia E, Oertli JJ (1994). Transport of ascorbic and dehydroascorbic acids across protoplast and vacuole membranes isolated from barley (*Hordeum vulgare* L. cv Gerbel) leaves. *Plant Physiol.* 106: 187-193.
- Reddy AR, Chaitanya KV, Munusamy-Vivekanandan (2004). Drought-induced responses of photosynthesis and antioxidant metabolism in higher plants. *J. Plant Physiol.* 161(11): 1189-1202.
- Saito K (1996). Formation of L-ascorbic acid and oxalic acid from D-glucosone in *Lemna minor*. *Phytochem.* 41: 145-149.
- Schwarz M (1985). The use of saline water in hydroponics. *Soilless Cult.* 1: 25-34.
- Smirnoff N (1995). Antioxidant systems and plant responses to the environment. In: Smirnoff N ed. *Environment and plant metabolism. Flexibility and acclimation.* Oxford: Bios Scientific Publishers, 217-243.
- Steel RGD, Torrie JH (1980). *Principles and procedures of statistics*, 2nd ed. McGraw-Hill Inc., USA.