

Full Length Research Paper

Impact of Fumonisin Purification on Immune Response in Experimental Paracoccidioidomycosis

A. A. Sasaki¹, E. Y. Hirooka², E. Y. S. Ono³, M. S. Kaminami¹, Pinge Filho P.¹, S. A. Khan¹, O. Kawamura⁴, A. B. Ribeiro², S. Fujii² and E. N. Itano^{1*}

¹Department of Pathological Science, State University of Londrina, P. O. Box 6001, 86051-990, Londrina, Paraná, Brazil.

²Department of Food Science and Technology, State University of Londrina, P. O. Box 6001, 86051-990, Londrina, Paraná, Brazil.

³Department of Biochemistry and Biotechnology, State University of Londrina, P. O. Box 6001, 86051-990, Londrina, Paraná, Brazil.

⁴Department of Biochemistry and Food Science, Kagawa University, Ikenobe, Miki-cho, Kita-gun, Kagawa, Japan.

Accepted 13 March, 2024

Fumonisin is a mycotoxin produced mainly by *Fusarium verticillioides*, which can modulate the immune response. Paracoccidioidomycosis (PCM), caused by the fungus *Paracoccidioides brasiliensis* (Pb), is one of the most important systemic mycoses in Latin America. The aim of this study was to evaluate the effect of the partially purified fumonisins on cellular immune response in mice infected with Pb. Four groups of male BALB/c mice were used. Groups PB and PB/FB were inoculated i.v. with 1×10^5 Pb yeast cells and, after 28 days, groups FB and PB/FB were inoculated (s.c.) with partially purified fumonisin B₁ from *F. verticillioides* (5×2.25 mg FB1/kg body weight). After 7 days, cellular immune response was evaluated by delayed-type hypersensitivity (DTH) and lymphoproliferative assays (LA) using spleen cells. Nitric oxide (NO) production by spleen cells was also evaluated. The specific LA response to Pb antigen was higher in group PB than in FB and CTR groups ($p < 0.05$) but not significant with PB/FB. The DTH response was higher in infected than non infected groups ($p < 0.05$) but also no significantly with PB and PB/FB groups. The lymphoproliferative response to ConA was decreased in FB or PB/FB in relation to CTR ($p < 0.05$) but not with PB/FB and also a reduction of NO levels was observed in fumonisin treated in relation to control group FB1/kg ($p < 0.05$). In conclusion, fumonisin B₁ or other components of *F. verticillioides* extracts significantly suppress the unspecific cellular immune response and the NO production by splenocytes from *P. brasiliensis* infected or not infected BALB/c mice.

Key words: Fumonisin, *Paracoccidioides brasiliensis*, lymphoproliferative assay, nitric oxide.

INTRODUCTION

Fumonisin is a group of mycotoxins produced by the plant pathogen *Fusarium verticillioides* and found predominantly in maize (Shephard et al., 1996). The extent of

contamination of raw corn with fumonisins varies with geographic location, agricultural and storage practices and the vulnerability of the plants to fungal invasion

*Corresponding author. Email: itanoeiko@hotmail.com.

during all phases of growth, storage and processing (Shelby et al., 1994). Brazil produces and consumes very large quantities of maize in a variety of forms, and Ono et al. (2000, 2001) reported fumonisin contamination in 98% of analyzed samples. Paracoccidioidomycosis (PCM) is a granulomatous disease caused by the thermally dimorphic fungus *Paracoccidioides brasiliensis* (Pb). The mycosis is endemic in Latin America and the majority of cases are reported in Brazil, where it is the 8th commonest cause of mortality among infectious diseases (Coutinho et al., 2002). PCM affects preferentially male farm workers (Coutinho et al., 2002; Franco et al., 1989). Airborne fungal propagules, consisting of conidia or hyphal fragments, begin the infection and undergo conversion to the yeast phase, the infective stage of PCM, in the lungs, progressing to haematogenic or lymphatic dissemination to the liver, spleen, skin and mucosa (McEwen et al., 1987). The disease may be developed in multiple forms, ranging from benign and localized (unifocal) to severe and disseminated (multifocal), depending on the host immune response (Marques et al., 2002). Fumonisin B₁ (FB₁) and their analogs are secondary toxic metabolites produced, mainly by *F. verticillioides* and *F. proliferatum* in maize (Leslie, 1996). The ingestion of FB₁ is related to many pathologies in humans as well as mice, rats, equines and swines (Sydenham et al., 1990; Ueno et al., 1997; Howard et al., 2001; Gelderblom et al., 1991; Marasas et al., 1988; Colvin and Harrison, 1992; Vizcarra-Olvera et al., 2012, Khan et al., 2012). The FB₁ can modulate the cellular or humoral immune response, affecting the antigen presenting cells, B and T lymphocytes and macrophages (Liu et al., 2002; Taranu et al., 2006; Martinova and Merrill, 1995; Devriendt, et al., 2009, Grenier et al., 2011, Bracarense et al., 2012, Burel et al., 2013). Fumonisin B₁ (FB₁) and their analogs are secondary toxic metabolites produced, mainly by *F. verticillioides* and *F. proliferatum* in maize (Leslie, 1996). The ingestion of FB₁ is related to much pathology in humans as well as mice, rats, equines and swines (Sydenham et al., 1990; Ueno et al., 1997; Howard et al., 2001; Gelderblom et al., 1991; Marasas et al., 1988; Colvin and Harrison, 1992). The FB₁ can modulate the cellular or humoral immune response, affecting the B and T lymphocytes activities and can also disrupt effector function of macrophages (Liu et al., 2002; Taranu et al., 2006; Martinova and Merrill, 1995). Due to chronicity of PCM and its prevalence in Brazil, there is a possibility of the concomitant contamination of fumonisin and Pb, mainly, maize is used as a basic feeding for the local poor community who are involved greatly in agriculture and the illness also affects the agricultural and poor population. In a previous experimental study, it was observed significantly increased the specific antibody response in male Swiss mice infected with Pb by FB₁ or other components of *F. verticillioides* extracts (Itano et al., 2008).

Considering the cellular immune response as the main

defense mechanism against Pb (Silva et al., 1995). The aim of this study was to evaluate the effect of the partially purified fumonisins on cellular immune response in mice infected with Pb-infected BALB/c mice.

MATERIALS AND METHODS

Fumonisin production and determination

F. verticillioides 103F (isolated from feed involved in animal intoxication in 1991 at Paraná State, Brazil) was grown on potato dextrose agar (PDA) slants at 25°C for 7 days. An aliquot of conidia suspension (10^6 conidia mL⁻¹) was inoculated into three 500 ml Erlenmeyer flasks containing 100 g ground maize adjusted to 42% moisture content and autoclaved twice for 30 min. The culture was incubated at 25°C for 14 days, extracted twice by shaking with ethyl acetate (1 L/kg) and filtered. Next, the residue was extracted once with 1 L/kg methanol: water (3:1, v/v) and twice with 1.5 L/kg of the same solvent. The dried extract was dissolved in 200 ml methanol: water (1:3, v/v) at 50°C and partitioned three times with 100 ml chloroform. The aqueous phase was dried at 50°C under vacuum (Cawood et al., 1991). This residue was dissolved in 20 ml methanol: water (1:3, v/v), and partially purified by passing successively through columns of Amberlite XAD-2 (100 g, 2.0 × 60 cm, Supelco, 10357-EA), silica gel (120 g, 2.0 × 60 cm, Sigma, 70 to 230 mesh, S2509), silica gel (120 g, 2.0 × 60 cm, Sigma, 70 to 230 mesh, S2509) and reversed-phase C18 (50 g, 3.0 × 30 cm, Acros, S2425) columns, using methanol, chloroform: methanol: acetic acid (6:3:1, v/v/v), chloroform: methanol: water: acetic acid (55:36:8:1, v/v/v/v) and a linear gradient from methanol: water (1:1, v/v) to methanol: water (4:1, v/v) as the mobile phases, respectively.

Fumonisin were determined by high-performance liquid chromatography (HPLC) (Shephard et al., 1990) with some modifications (Ueno et al., 1993). An aliquot of partially purified extract (6 ml) was filtered through a 0.45-µm syringe filter (Millipore, Windham, US) and suitably diluted with acetonitrile: water (1:1, v/v). After derivatization with 200-µL of o-phthalaldehyde (OPA) reagent, the sample was injected into the HPLC system within 1 min. Fumonisin were analyzed by a reversed-phase, isocratic system (Shimadzu LC-10 AD pump and RF-10A XL fluorescence detector), using a Shim-pack CLC-ODS (M) column (4.6 × 250 mm, Shimadzu). Fluorescence excitation and emission wavelengths were 335 and 450 nm, respectively. The eluent was CH₃OH: 0.1 M NaH₂PO₄ (80:20, v/v), adjusted to pH 3.3 with H₃PO₄ solution, flowing at 1 ml/min. The detection limits for the fumonisins FB₁ and FB₂ were 27.5 and 35.3 ng/g, respectively. FB₁ and FB₂ levels in the partially purified extract were 152.1 and 53.34 mg/mL, respectively. The sample was diluted in sterile PBS to a concentration of 22.5 mg FB₁/ml, this partially purified fumonisin B₁ (PPF) was used for animal treating.

P. brasiliensis

P. brasiliensis strain 18 (Pb 18) was cultured on Sabouraud agar (Micromed, Rio de Janeiro, RJ, Brazil), and maintained at 35°C, subculturing at 5 day intervals. The yeast cells were collected in sterile saline, filtered through cotton and gauze layers, and the concentration was adjusted to 1×10^6 cells/ml.

Experimental design

Male BALB/c mice (n = 24), 4 to 5 weeks old (20 to 25 g), provided by the animal facilities of Londrina State University (UEL), were kept in a 12:12 h light-dark cycle at 25°C, with free access to sterilized commercial mouse food (Nuvital, Curitiba, Brazil) and water. Four groups of 6 mice were used: infected (PB), treated (FB),

infected and treated (PB/FB) and uninfected and untreated as control (CTR). Groups PB and PB/FB were inoculated (i.v.) with 1×10^5 yeast cells (Pb 18) and, 28 days later, groups FB and PB/FB were inoculated (s.c.) with PPF from *F. verticillioides* (5×2.25 mg FB1/kg body weight/day), as described by Johnson and Sharma (2001). The control group was inoculated with sterile PBS. Plasma samples and spleens cells were taken 7 days after PPF inoculation. The procedures and experimental design were approved by the Animal Research Ethics Committee of the UEL.

Exoantigen (exoAg)

A lyophilized exoAg was prepared from a yeast-phase culture of Pb strain B-339 according to Camargo et al. (1989).

Lymphoproliferation assay

Spleen cells from infected and normal mice were removed aseptically and teased. The erythrocytes were lysed with tris-ammonium chloride, and the cell suspension was washed three times in RPMI medium. For lymphoproliferative assays, 100 μ l of the cells were cultured in triplicate wells at a concentration of 1×10^6 cells/ml in RPMI 1640 containing L-glutamine (Sigma-Aldrich, St. Louis, Missouri, USA), 10% fetal calf serum (Gibco, USA) in 96-well flat-bottom culture plates. ExoAg or concanavalin A (Con) (Sigma-Aldrich, St. Louis, Missouri, USA) was then added to each well at concentrations of 50 and 10 μ g/ml, respectively and as control with RPMI medium. The cells were cultured for 3 (ConA) or 5 (exoAg) days at 37°C with 5% CO₂ and were pulsed with 1 μ Ci of (³H) thymidine 18 h before harvesting on glass filter strips. The radioactivity was determined by a liquid scintillation counter (Beckman LS 6.800). The 'stimulation index' was calculated as the triplicate of stimulated cells (cells + ExoAg or ConA) divided by the cell control (cell + RPMI).

Delayed-type hypersensitivity assay (DTH)

The DTH reactions were performed in all groups of animals. Mice were inoculated with exoAg (10 μ g) and sterile PBS in right and left footpad respectively. The footpad thickness was measured at 24 h after inoculation with caliper (Mitotoyo, Tokyo, Japan). The increase in thickness was calculated, as in the formula ($A = D^2/4, = 3.14$).

Measurement of nitric oxide (NO)

The concentration of NO in spleens cells was measured by a microplate Griess assay. Spleen cells (100 μ L) prepared as described earlier, were cultured with lipopolysaccharides (LPS), (7 g) during 24 h at 37°C. The supernatants were collected and 50 μ L of these samples were transferred to wells of 96-well flat bottom microtiter plate and incubated with an equal volume of the Griess reagent for 10 min at room temperature. The absorbance at 550 nm was determined with a Titertek Multiskan apparatus (Multiskan EX, Uniscience – Labsystems, Helsinki, Finland). The NO concentration was determined using a standard curve determined with different concentrations of NaNO₂.

Statistical analysis

The data were analysed by using a Tukey-Kramer ANOVA test for comparisons and declaring significance at $p < 0.05$.

RESULTS

Proliferation response of spleens cells to exoAg and to ConA

During the present investigation, higher proliferation response to exoAg in infected groups (PB: 2.04 ± 0.71 ; PB/FB: 1.55 ± 0.26) in relation to not infected (FB: 0.87 ± 0.15 ; CTR; 1.10 ± 0.23) was observed (Figure 1). The decreased proliferation response in PB/FB was not statistically significant in relation to PB group. CTR \times PB ($p < 0.01$), FB \times PB ($p < 0.001$) and FB \times PB/FB ($p < 0.05$). Figure 2 shows decreased proliferation response to ConA in FB treated groups, both in infected or not infected (FB: 0.12 ± 0.08 and PB/FB: 0.17 ± 0.24) in relation to not treated (PB: 1.52 ± 0.28 and CTR: 1.95 ± 0.41). CTR \times FB and PB/FB, $p < 0.001$.

DTH assay

Results regarding the 'delayed type hypersensitivity' in the infected groups showed higher inflammatory response than control not infected group (PB: 1.59 ± 0.19 ; PB/FB: 1.49 ± 0.09 , FB: 0.98 ± 0.08 ; CTR: 1.10 ± 0.16); $p < 0.001$, but these results are similar between PB and PB/FB, $p > 0.05$ (Figure 3).

NO concentration in culture supernatants of spleens cells stimulated with LPS

During the present study, 'nitrous oxide' amount was measured in the supernatants of spleens stimulated with LPS and it was detected decreased level in FB treated or infected with Pb or infected and treated with PPF in relation to control group ($p < 0.001$), but similar between FB treated or infected with Pb or infected and treated with PPF ($p > 0.05$) (Figure 4).

DISCUSSION

In PCM, the cellular immune response represents the main defense mechanism against Pb and in this study, it was observed an induction of specific cellular response in infected mice both by DTH and LA assays. However, it was not observed significant suppression in simultaneous infection with Pb and contaminated with PPF from *F. verticillioides* in experimental PCM in male BALB/c mice. However, there were a significant suppression of response to ConA in mice treated with PPF in relation to the normal control without treatment with PPF, suggesting that the FB₁ in our conditions of experiment does not affect the specific cellular immune response but can suppress in specific immune response. Our results are partially in accordance with the data of Theumer et al. (2002), which




Figure 1. Proliferation response of spleens cells to exoAg. Spleen cells (1×10^5 cells/ml) were cultured with *P. brasiliensis* exoAg (50 $\mu\text{g/ml}$) for 5 days, pulsed with 1 μCi and the stimulation index calculated as the triplicate of stimulated cells divided by the control not stimulated cells. A higher proliferation response to exoAg was observed in infected groups in relation to not infected. PB/FB = BALB/c mice infected with *P. brasiliensis* (1×10^5 yeast cells, 28 days) and treated with PPF from *F. verticillioides* (5×2.25 mg FB₁/kg body weight/day); PB = only infected; FB = only treated with PPF and CTR = inoculated with sterile PBS. Different letters indicate significant differences between groups ($p < 0.05$).




Figure 2. Stimulation of cellular proliferation with ConA. Spleen cells (1×10^5 cells) were cultured with ConA (10 $\mu\text{g/ml}$) for 72 h, pulsed with 1 μCi and the stimulation index (SI) calculated as the triplicate of stimulated cells divided by the control not stimulated cells. PB/FB = BALB/c mice infected with *P. brasiliensis* (1×10^5 yeast cells, 28 days) and treated with PPF from *F. verticillioides* (5×2.25 mg FB₁/kg body weight/day); PB = only infected; FB = only treated with PPF and CTR = inoculated with sterile PBS. SI were more elevated in CTR and Pb groups than PBS/FB and PB/FB groups; however, significant results were found only between CTR and PBS/FB, and between PBS and Pb/FB. Different letters indicate significant differences between groups ($p < 0.05$).




Figure 3. DTH assay. BALB/c mice were infected with *P. brasiliensis* (1×10^5 yeast cells, 28 days) and treated with PPF from *F. verticillioides* (5×2.25 mg FB1/kg body weight/day) (PB/FB) or only infected (PB) or only treated with PPF (FB) or inoculated with sterile PBS (CTR). ExoAg from *P. brasiliensis* were injected intra-footpad 24 h before measurement of the footpad response. Different letters indicate significant differences between groups ($p < 0.05$).




Figure 4. NO determination. Spleen cells were stimulated with LPS to determine the release of NO in the culture medium. PB/FB = BALB/c mice infected with *P. brasiliensis* (1×10^5 yeast cells, 28 days) and treated with PPF from *F. verticillioides* (5×2.25 mg FB1/kg body weight/day); PB = only infected; FB = only treated with PPF and CTR = inoculated with sterile PBS. Inhibition of NO was seen in groups infected and treated with PPF. Different letters indicate significant differences between groups ($p < 0.05$).

they had observed no significant effect on mitogen-induced proliferation of spleen cells in mice treated with FB₁ and stimulated with phytohemagglutinin mitogen (PHA). The inflammatory reaction is a non-specific response and Gonzalez et al. (2000) reported that the NO produced by spleen macrophages has crucial role in *P. brasiliensis*

defense. Our NO analysis demonstrated decreased levels when spleen macrophages from groups of infected or PPF treated or infected/treated mice stimulated with LPS in relation to control not treated and not infected group, suggesting that both PPF and Pb can modulate the NO production by spleen macrophages, but not with synergic

effect. Also, Popi et al. (2004) reported the decreased NO production by peritoneal macrophages incubated with gp43 and stimulated with zymozan.

The infection was confirmed by histologic analysis of lungs and liver, where no significant difference between the infected and infected/treated group was observed (data not shown), suggesting that in the conditions of the experiment the mycotoxin did not affect the gravity of the illness. All groups of animals had the same initial healthy appearance, which they maintained throughout the period of the study. This finding is in accordance with the results of evaluation of the specific cellular immune response, the main protective response against the Pb (Silva et al., 1995; Calich and Kashino, 1998; Kashino et al., 2000). In Bhandari et al. (2001) work, it was reported that FB₁ modulates the IFN- γ , a Th1 marker cytokine in BALB/c mice according to the sex, with elevated expression (male) or down-modulation (female). The lack of change in specific cellular immune response in BALB/c mice (male) infected and treated with partially purified fumonisin could be due to the balance between FB₁ (Bhandari et al., 2001) and immunosuppressive Pb antigens actions (Benard et al., 1997; Cacere et al., 2002 and Rigobello et al., 2013), that require further study.

The literature data has shown the low proliferation of spleenocytes to some Pb antigens (Benard et al., 1997), that induce apoptosis and suggested that it may be one of the mechanisms leading to hyporesponsiveness (Cacere et al., 2002). The relevance of this study stems from the fact that PCM is the 8th most common cause of death from predominantly chronic or recurring infectious and parasitic diseases in Brazil. It is believed that about 10 million people are infected with the fungus, and 2% of them may develop the disease (McEwen et al., 1995). Thus, any factor that leads to modulation of the immune response, such as the presence of mycotoxin, could contribute to the development of the disease, increase its severity or control the disease. For the best understanding about the action of the FB₁ in the course of the experimental PCM, there are necessary additional studies using purified FB₁ in diverse concentrations/times of incubation/mouse isogenic lines by using both sex. We concluded for the results that FB or other components of *F. verticillioides* extracts does not affect specific cellular immune response in experimental PCM in male BALB/c mice, but can affect non-specific response, in the conditions of the work.

ACKNOWLEDGEMENTS

This work was supported by CAPES, CNPq, Fundação Araucária, FAEP/UEL. The authors thank Timothy J.C. Roberts for the careful correction of the English text.

REFERENCES

Benard G, Mendes-Giannini MJ, Juvenale M, Miranda ET, Duarte AJ (1997). Immunosuppression in paracoccidioidomycosis: T cell

- hyporesponsiveness to two *Paracoccidioides brasiliensis* glycoproteins that elicit strong humoral immune response. *J. Infect. Dis.* 175:1263-1267.
- Bhandari N, He Q, Sharma RP (2001). Gender-related differences in subacute fumonisin B1 hepatotoxicity in BALB/c mice. *Toxicol.* 165:195-204.
- Bracarense AP, Luciola J, Grenier B, Drociunas Pacheco G, Moll WD, Schatzmayr G, Oswald IP (2012). Chronic ingestion of deoxynivalenol and fumonisin, alone or in interaction, induces morphological and immunological changes in the intestine of piglets. *Br. J. Nutr.* 107(12):1776-86.
- Burel C, Tanguy M, Guerre P, Boilletot E, Cariolet R, Queguiner M, Gilbert Postollec, Pinton P, Salvat G, Oswald IP, Fravallo P (2013). Effect of Low Dose of Fumonisin on Pig Health: Immune Status, Intestinal Microbiota and Sensitivity to *Salmonella*. *Toxins (Basel)*. 5(4): 841-864.
- Cacere CR, Romano CC, Mendes Giannini MJ, Duarte AJ, Benard G (2002). The role of apoptosis in the antigen-specific T cell hyporesponsiveness of paracoccidioidomycosis patients. *Clin. Immunol.* 105:215-222.
- Calich VL, Kashino SS (1998). Cytokines produced by susceptible and resistant mice in the course of *Paracoccidioides brasiliensis* infection. *Braz. J. Med. Biol. Res.* 31:615-623.
- Camargo ZP, Unterkircher C, Travassos LR (1989). Identification of antigenic polypeptides of *Paracoccidioides brasiliensis* by immunoblotting. *J. Med. Vet. Mycol.* 27:407-412.
- Cawood ME, Gelderblom WCA, Vleggaar R, Behrend Y, Thiel PG et al (1991). Isolation of the fumonisin mycotoxins: a quantitative approach. *J. Agric. Food Chem.* 39:1958-1962.
- Colvin BM, Harrison LR (1992). Fumonisin-induced pulmonary edema and hydrothorax in swine. *Mycopathology* 117:79-82.
- Coutinho ZF, Silva D, Lazera M, Petri V, Oliveira RM et al (2002). Paracoccidioidomycosis mortality in Brazil (1980-1995). *Cad Saude Publica.* 18:1441-1454.
- Devriendt B, Gallois M, Verdonck F, Wache Y, Bimczok D, Oswald IP, Goddeeris BM, Cox E (2009). Remove from marked Records The food contaminant fumonisin B₁ reduces the maturation of porcine CD11R1⁺ intestinal antigen presenting cells and antigen-specific immune responses, leading to a prolonged intestinal ETEC infection. *Veterinary Research* 40(4):40.
- Franco MF, Mendes RP, Rezkalla-Iwasso M, Montenegro MR (1989). Paracoccidioidomycosis. *Baillière's Clin. Trop. Medicine Commu. Dis.* 4:185-219.
- Gelderblom WC, Kriek NP, Marasas WF, Thiel PG (1991) Toxicity and carcinogenicity of the *Fusarium moniliforme* metabolite, fumonisin B₁ in rats. *Carcinog.* 12:1247-1251.
- Gonzalez A, de Gregori W, Velez D, Restrepo A, Cano LE (2000). Nitric oxide participation in the fungicidal mechanism of gamma interferon-activated murine macrophages against *Paracoccidioides brasiliensis* conidia. *Infect. Immun.* 68:2546-2552.
- Grenier B, Loureiro-Bracarense AP, Luciola J, Pacheco GD, Cossalter AM, Moll WD, Schatzmayr G, Oswald IP (2011). Individual and combined effects of subclinical doses of deoxynivalenol and fumonisins in piglets. *Mol Nutr Food Res.* 55(5):761-771.
- Howard PC, Eppley RM, Stack ME, Warbritton A, Voss KA et al (2001). Fumonisin b1 carcinogenicity in a two-year feeding study using F344 rats and B6C3F1 mice. *Environ. Health Perspect.* 109 Suppl. 2:277-282.
- Itano EN, Sasaki AA, Ribeiro AB, Fujii S, Ono EY et al (2008). Effect of *Fusarium verticillioides* Extract on Specific Antibody Production Against *Paracoccidioides brasiliensis*. *World Mycotoxin J.* 1:375-380.
- Johnson VJ, Sharma RP (2001). Gender-dependent immunosuppression following subacute exposure to fumonisin B1. *Int. Immunopharmacol.* 1:2023-2034.
- Kashino SS, Fazioli RA, Cafalli-Favati C, Meloni-Bruneri LH, Vaz CA et al (2000). Resistance to *Paracoccidioides brasiliensis* infection is linked to a preferential Th1 immune response, whereas susceptibility is associated with absence of IFN- γ production. *J. Interferon. Cytokine Res.* 20:89-97.
- Khan MA, Iqbal A, Asrani RK (2012) Microscopic changes due to fumonisin B and ochratoxin A 1 induced nephropathy in Japanese

- quail. *Vet. World*, 5(9): 535-540.
- Leslie JF (1996). Introductory biology of *Fusarium moniliforme*. *Adv. Exp. Med. Biol.* 392:153-164.
- Liu BH, Yu FY, Chan MH, Yang YL (2002). The effects of mycotoxins, fumonisin B₁ and aflatoxin B₁, on primary swine alveolar macrophages. *Toxicol. Appl. Pharmacol.* 180:197-204.
- Marasas WF, Kellerman TS, Gelderblom WC, Coetzer JA, Thiel PG et al (1988). Leukoencephalomalacia in a horse induced by fumonisin B₁ isolated from *Fusarium moniliforme*. *Onderstepoort J. Vet. Res.* 55:197-203.
- Marques Mello L, Silva-Vergara ML, Rodrigues V Jr (2002) Patients with active infection with *Paracoccidioides brasiliensis* present a Th2 immune response characterized by high Interleukin-4 and Interleukin-5 production. *Hum. Immunol.* 63:149-154.
- Martinova EA, Merrill AH Jr (1995) Fumonisin B₁ alters sphingolipid metabolism and immune function in BALB/c mice: immunological responses to fumonisin B₁. *Mycopathol.* 130:163-170.
- McEwen JG, Bedoya V, Patino MM, Salazar ME, Restrepo A (1987) Experimental murine paracoccidioidomycosis induced by the inhalation of conidia. *J. Med. Vet. Mycol.* 25:165-175.
- McEwen JG, Garcia AM, Ortiz BL, Botero S, Restrepo A (1995). In search of the natural habitat of *Paracoccidioides brasiliensis*. *Arch. Med. Res.* 26:305-306.
- Ono EY, Kawamura O, Ono MA, Ueno Y, Hirooka EY (2000). A comparative study of indirect competitive ELISA and HPLC for fumonisin detection in corn of the State of Paraná, Brazil. *Food Agric. Immunol.* 12:5-14.
- Ono EY, Ono MA, Funo FY, Medinal AE, Oliveira TC et al (2001). Evaluation of fumonisin-aflatoxin co-occurrence in Brazilian corn hybrids by ELISA. *Food Addit. Contam.* 18:719-729.
- Popi AF, Lopes JD, Mariano M (2004). Interleukin-10 secreted by B-1 cells modulates the phagocytic activity of murine macrophages in vitro. *Immunology.* 113:348-354.
- Rigobello FF, Marquez AS, Lopes JD, Nakanishi-Ito FA, Itano EN (2013). Patients with Chronic-form *Paracoccidioidomycosis* Present High Serum Levels of IgE Anti-*paracoccidioides brasiliensis* Gp70. *Mycopathologia* 175:307-313.
- Shelby RA, White DG, Bauske EM (1994). Differential fumonisin production in maize hybrids. *Plant Dis.* 78:582-584.
- Shephard GS, Sydenham EW, Thiel PG, Gelderblom WCA (1990). Quantitative determination of fumonisins B₁ and B₂ by high-performance liquid chromatograph with fluorescence detection. *J. Liq. Chromatogr.* 13:2077-2087.
- Shephard GS, Thiel PG, Stockenstrom S, Sydenham EW (1996). Worldwide survey of fumonisin contamination of corn and corn-based products. *J. AOAC Int.* 79:671-687.
- Silva CL, Silva MF, Faccioli LH, Pietro RC, Cortez SA et al (1995). Differential correlation between interleukin patterns in disseminated and chronic human paracoccidioidomycosis. *Clin. Exp. Immunol.* 101:314-320.
- Sydenham EW, Thiel PG, Marasas WFO, Shephard GS, Van Schalkwyk DJ et al (1990). Natural occurrence of some *Fusarium* mycotoxins in corn from low and high esophageal cancer prevalence areas of the Transkei, South Africa. *J. Agric. Food Chem.* 38:1900-1903.
- Taranu I, Marin DE, Bouhet S, Pascale E, Bailly JD et al (2006). *In vivo* and *in vitro* effects of fumonisin B₁ on cytokine production: implication for the immune response to vaccination. *Mycotoxins and Phycotoxins* 1:157-167.
- Theumer MG, Lopez AG, Masih DT, Chulze SN, Rubinstein HR (2002). Immunobiological effects of fumonisin B₁ in experimental subchronic mycotoxicoses in rats. *Clin. Diagn. Lab. Immunol.* 9:149-155.
- Ueno Y, Ayoama S, Sigiura Y, Wang D-S, Hirooka EY et al (1993). A limited survey of fumonisin in corn and corn-based products in Asian countries. *Mycotoxin Res.* 9:27-34.
- Ueno Y, Iijima K, Wang SD, Sugiura Y, Sekijima M et al (1997). Fumonisin as a possible contributory risk factor for primary liver cancer: a 3-year study of corn harvested in Haimen, China, by HPLC and ELISA. *Food Chem. Toxicol.* 35:1143-1150.
- Vizcarra-Olvera JE, Astiazarán-García H, Burgos-Hernández A, Parra-Vergara NV, Cinco-Moroyoqui FJ, Sánchez-Mariñez RI, Quintana-Obregón EA, Cortez-Rocha MO (2012). Evaluation of pathological effects in broilers during fumonisins and clays exposure. *Mycopathologia.* 174(3):247-54.