

## Full Length Research Paper

# Sunflower seed invigoration by chemical manipulation

Chandan Kumar Pati<sup>1\*</sup> and Alope Bhattacharjee<sup>2</sup><sup>1</sup>Department of Botany, Saldiha College, Saldiha-722 173, Bankura, West Bengal, India.<sup>2</sup>UGC Centre for Advanced Study, Department of Botany, University of Burdwan, Burdwan-713 104, West Bengal, India.

Accepted 13 April, 2019

Sunflower (*Helianthus annuus* L. cv. Morden) seeds lose viability at a rapid pace under accelerated ageing (AA) condition. Pretreatment of the seeds with salicylic acid and ascorbic acid for 8 h before imposition of accelerated ageing treatment in desiccator (99.1% RH and  $32 \pm 2^\circ\text{C}$ ) or continuous treatment of the seeds with *Eucalyptus* oil vapour for 60 days in a separate desiccator under the same ageing condition slowed down the ageing-induced rapid loss of germination. Plant performance was found to be much better when they were developed from seeds which underwent chemical pretreatment measured in terms of field emergence capacity, root length, shoot length, fresh weight and dry weight of plants. Plant potential was also higher in the pretreatments as evidenced from the treatment-induced higher chlorophyll, protein, DNA and RNA levels as well as activities of catalase and peroxidase enzymes in spite of adverse storage situation. Yield attributes like diameter and fresh weight of capitulum, seed number per capitulum, 1000 seed weight were found significantly higher in plants raised from pretreated seed samples. Results, therefore, pointed out that in spite of experiencing accelerated ageing treatment, the chemical-pretreated seeds retained higher seed vigor and produced healthier plants which resulted in enhancement of crop yield.

**Key words:** Eucalyptus oil, accelerated ageing, ascorbic acid, salicylic acid, seed viability, sunflower, yield.

## INTRODUCTION

Storing of seeds is a serious problem in tropical and subtropical countries where high temperature and high relative humidity greatly accelerate seed ageing phenomenon causing consequent deterioration and non-viability of seeds. The problem of retention of seed vigor in many states of India is much more acute because of its semiarid climate where high relative humidity prevailing during the major part of a year is very conducive to the growth of microorganisms, particularly fungi (Christensen and Kaufmann, 1965; Halder, 1981; Aziz and Shahir, 1997). As most crop seeds require storage for either one or several planting seasons, agriculturists and horticulturists of this region are often handicapped with respect to maintenance of standard seed vigor under ambient storage environment. The problem becomes more acute as seeds experience drought and other stress periodically under storage. Seed vigor and viability are two interrelated characters and generally loss of vigor

precedes loss of viability. As per the principles proposed by Copland and McDonald (2001), Rooszrokh et al. (2002), Sengupta et al. (2005) and Barman (2006), we may look upon vigor as the quality of the seed responsible for rapid and uniform germination, extended storability and perform a good field emergence. Keeping this problem of seed storage in mind, an attempt was made in this investigation to prolong the storage life of a sunflower cultivar having viability problems. Present experiment was performed under accelerated ageing (AA) condition by imposing high relative humidity with a view to maintaining a uniform adverse storage condition and also to obtain expeditious results. In fact, accelerated ageing treatment provides a powerful manipulative tool which makes it possible to study the process of seed deterioration over a very short period (Heydecker, 1972) and this mimics the natural ageing process (Delouche and Baskin, 1973; Perl et al., 1978; Halder, 1981; Pati and Bhattacharjee, 2011).

Although efficacy of several classes of chemicals namely hormones, retardants, redox chemicals, phenols, vitamins and salts on maintenance of seed health under

\*Corresponding author. E-mail [cpbotany@yahoo.co.in](mailto:cpbotany@yahoo.co.in).

**Table 1.** Effect of seed pretreatment with salicylic acid (S.A., 200 g/ml) and ascorbic acid (A. A., 200 g/ml) and treatment with *Eucalyptus* oil (*E. oil*) on germination and field emergence capacity of sunflower seeds stored under AA condition for 60 days.

Treatment	Percentage germination				Field emergence capacity (%)			
	AA (days)							
	0	20	40	60	0	20	40	60
Control	100	48.6 <sup>a</sup>	27.5 <sup>a</sup>	NA	87.9 <sup>a</sup>	38.5 <sup>c</sup>	12.4 <sup>a</sup>	NA
S.A.	100	56.6 <sup>c</sup>	39.9 <sup>c</sup>	NA	90.2 <sup>a</sup>	40.1 <sup>c</sup>	18.0 <sup>c</sup>	NA
A.A.	100	54.4 <sup>bc</sup>	35.2 <sup>b</sup>	NA	91.2 <sup>a</sup>	36.6 <sup>b</sup>	14.1 <sup>b</sup>	NA
E.oil	100	51.8 <sup>b</sup>	33.5 <sup>b</sup>	NA	88.5 <sup>a</sup>	31.5 <sup>a</sup>	12.3 <sup>a</sup>	NA
LSD(P=0.05)	NC	2.05	1.81	-	NS	2.90	1.20	-

Seeds were presoaked with salicylic acid, ascorbic acid or distilled water for 8 h and then sun dried, or seeds were given continuous treatment with the vapour of *Eucalyptus* oil. The seed samples were then separately allowed to experience AA treatment (99.1% RH) in a desiccator. Data were recorded after 0, 20, 40 and 60 days of AA seeds. Values in a vertical column followed by the same letter are not significantly different at 5% level. NA: No germination, NC: Not calculated, NS: Not significant.

storage has been established (Bhattacharyya and Basu, 1990; Chhetri et al., 1993; Basu, 1994; Rai, 2000, this field of seed physiology still remains relatively less explored. Thus, the major objective of this work was to test the efficacy of a growth retardant (salicylic acid), an antioxidant (ascorbic acid) and a volatile oil (*Eucalyptus* oil) on the alleviation of deterioration and viability extension of sunflower seeds under storage.

## MATERIALS AND METHODS

After surface sterilization (0.1% HgCl<sub>2</sub> for 90 s) certified hybrid seeds of sunflower (*Helianthus annuus* L. cv. Morden) were separately presoaked in the different concentration grades of aqueous solutions of salicylic acid, ascorbic acid or distilled water for different durations. And the seeds were then dried back to original dry weight by intense sun drying followed by measuring of the moisture level. After initial screening to allow maximum penetration of the experimental chemicals, it was found that the concentrations of salicylic acid (200 g/ml) and ascorbic acid (200 g/ml) at 8 h were more effective to the present experiment. The pretreated seed lots (200 g each) were taken in separate porous cloth bags and thus stored in a desiccator in which 99.1% relative humidity (RH) was preimposed by keeping 250 ml 3.03% H<sub>2</sub>SO<sub>4</sub> within it (Rao et al. 2003). This experimental set-up was kept at 32 ± 2°C for 60 days allowing the seeds to experience forced ageing treatment and H<sub>2</sub>SO<sub>4</sub> was changed at 7-day intervals to restore the desired RH within the desiccator for 60 days.

In a separate experiment, a seed lot (200 g) was kept in a smaller desiccator in which 5 ml *Eucalyptus* oil extracted from mature leaves, procured from Shri Mahavir Eucalyptus Oil Company, Tamil Nadu, India was taken in a small Petri dish in addition to 250 ml 3.03% H<sub>2</sub>SO<sub>4</sub>. Here the seeds underwent treatment with the vapour of *Eucalyptus* oil along with accelerated ageing treatment (99.1% RH) throughout the experimental period. From the seed lots of both the experiment, germination and field emergence capacity of seeds were made after 0, 20, 40 and 60 days of accelerated ageing treatment.

To analyse the percentage germination, four groups of 100 seeds (total 400 seeds) were transferred to separate Petri dishes containing filter paper moistened with distilled water. Percentage germination data were recorded after 168 h of seed soaking

following the International Rules for Seed Testing (ISTA, 1976). Percentage field emergence capacity were recorded after emerging of radicle and plumule from the treated seed samples sown in the experimental field after 10 days of seed sowing (Bhattacharjee, 1984; Pati, 2007).

Some growth and biochemical parameters as well as yield attributes were recorded from the plants raised from the accelerated aged (0 and 20 days only) seeds. Plants were established and data were recorded at two developmental stages viz. pre-heading stage (30-days-old plants) and post-heading stage (60-days-old plants) from 10 uniformly grown plants of each treatment.

Extraction and estimation of chlorophyll and protein from mature leaves of each treated samples were done by the method of Arnon (1949) and Lowry et al. (1951) respectively. Extraction of nucleic acids was made from the same leaves following the method described by Cherry (1962) and quantitative estimation was done as per the method of Markham (1955) modified by Choudhuri and Chatterjee (1970). Activity of catalase (CAT) was analysed following the method of Snell and Snell (1971) as modified by Biswas and Choudhuri (1978) and that of peroxidase (POD) was analysed as per the method of Kar and Mishra (1976). For assaying of the enzymes, the blank was taken as zero time control and the activity was expressed as  $(\Delta A \times T_v) / (t \times v)$ , where  $\Delta A$  is the absorbance of the sample after incubation minus the absorbance of the zero time control,  $T_v$  is the total volume of the filtrate,  $t$  is the time (minutes) of incubation with the substrate and  $v$  is the volume of the filtrate taken for incubation (Fick and Qualset, 1975).

Yield attributes recorded include: diameter and fresh weight of capitulum, seed number per capitulum and 1000 seed weight. Statistical analysis of the data was done in terms of least significant difference (LSD) which was calculated at 95% confidence limits (Panse and Sukhatme, 1967).

## RESULTS AND DISCUSSION

Results showed that pretreatment of sunflower seeds with salicylic acid, ascorbic acid and *Eucalyptus* oil significantly alleviated the ageing-induced loss of germination and enhanced field emergence capacity under accelerated ageing environment (Table 1). Reduced seed germinability and field emergence capacity are considered to be the important visible criteria

**Table 2.** Effect of seed pretreatment with salicylic acid (S.A., 200 µg/ml) and ascorbic acid (A. A., 200 µg/ml) and treatment with *Eucalyptus* oil (E. oil) followed by AA treatment for 20 days on changes of root length and shoot length of sunflower plants.

Treatment	Root length (cm)				Shoot length (cm)			
	Plant age (days)							
	30		60		30		60	
	AA(days)							
	0	20	0	20	0	20	0	20
Control	5.09 <sup>a</sup>	4.23 <sup>a</sup>	14.18 <sup>a</sup>	10.01 <sup>a</sup>	28.9 <sup>b</sup>	12.0 <sup>a</sup>	82.5 <sup>a</sup>	69.5 <sup>a</sup>
S.A.	5.36 <sup>a</sup>	5.12 <sup>c</sup>	15.68 <sup>bc</sup>	12.83 <sup>b</sup>	25.2 <sup>a</sup>	15.3 <sup>c</sup>	100.2 <sup>c</sup>	83.5 <sup>cd</sup>
A.A.	5.20 <sup>a</sup>	5.08 <sup>c</sup>	16.89 <sup>c</sup>	12.99 <sup>b</sup>	32.4 <sup>c</sup>	18.2 <sup>d</sup>	110.6 <sup>d</sup>	87.5 <sup>d</sup>
E.oil	5.13 <sup>a</sup>	4.95 <sup>bc</sup>	15.40 <sup>bc</sup>	12.56 <sup>b</sup>	30.5 <sup>bc</sup>	13.3 <sup>b</sup>	91.2 <sup>b</sup>	76.8 <sup>bc</sup>
LSD(P=0.05)	0.41	0.33	1.31	1.01	1.90	1.22	8.21	6.85

Data were recorded from 30 and 60 days old uniformly grown plants. Values in a vertical column followed by the same letter are not significantly different at 5% level.

**Table 3.** Effect of seed pretreatment with salicylic acid(S.A., 200 µg/ml) and ascorbic acid (A. A., 200 µg/ml) and treatment with *Eucalyptus* oil (E. oil) followed by AA treatment for 20 days on changes of fresh weight and dry weight of sunflower plants.

Treatment	Fresh weight (g)				Dry weight (g)			
	Plant age (days)							
	30		60		30		60	
	AA(days)							
	0	20	0	20	0	20	0	20
Control	36.0 <sup>a</sup>	32.5 <sup>a</sup>	398.9 <sup>a</sup>	321.0 <sup>a</sup>	12.7 <sup>a</sup>	7.5 <sup>a</sup>	52.4 <sup>a</sup>	41.9 <sup>a</sup>
S.A.	54.3 <sup>d</sup>	47.2 <sup>c</sup>	445.8 <sup>b</sup>	406.3 <sup>d</sup>	16.6 <sup>c</sup>	11.0 <sup>d</sup>	78.3 <sup>d</sup>	71.5 <sup>d</sup>
A.A.	51.1 <sup>cd</sup>	46.3 <sup>c</sup>	434.6 <sup>b</sup>	375.1 <sup>c</sup>	15.8 <sup>bc</sup>	10.1 <sup>c</sup>	72.1 <sup>c</sup>	67.1 <sup>c</sup>
E.oil	43.2 <sup>b</sup>	40.5 <sup>b</sup>	401.5 <sup>a</sup>	350.7 <sup>b</sup>	15.4 <sup>b</sup>	9.4 <sup>b</sup>	64.7 <sup>b</sup>	58.2 <sup>b</sup>
LSD(P=0.05)	3.55	3.22	26.85	22.08	1.05	0.60	5.26	4.20

Data were recorded from 30 and 60 days old uniformly grown plants. Values in a vertical column followed by the same letter are not significantly different at 5% level.

for the evaluation of poor seed vigor (Anderson, 1970, Halder et al., 1983; Rai, 2000). In this investigation, the chemical-induced cessation of loss of seed germination and field emergence capacity are indicative of storage potential enhancement property of the test chemicals.

Accelerated ageing treatment for 20 days impaired field performance of sunflower plants as evident from the reduction of root length and shoot length (Table 2), fresh weight and dry weight (Table 3), levels of chlorophyll and protein (Table 4), DNA and RNA (Table 5) as well as activities of CAT and POD enzymes (Table 6). The chemical-induced alleviation of the deleterious effects of ageing on the overall growth and metabolism of sunflower plants which indicates the effectiveness of pretreatment by salicylic acid, ascorbic acid and *Eucalyptus* oil. The ageing-induced adverse effects on overall growth and metabolism of sunflower plant were

also reflected in some yield attributes. Reduced field performance of plants was associated with concomitant reduction of yield attributes leading to impairment of final seed yield (Table 7) of the plants which were developed from the forced aged seeds. Here also, salicylic acid, ascorbic acid and *Eucalyptus* oil showed a promising role as the adverse effects on plant development and crop yield were alleviated to a considerable extent.

Chlorophyll, protein, DNA, RNA, CAT and POD are regarded as reliable indices of vigor status of plants. In this investigation, comparatively better plant health and higher metabolic status of plants, raised from the chemical-treated seeds, are indicative of invigoration of seeds under storage. And the invigorated seeds subsequently exhibited better field performance which was recorded in terms of plant growth and metabolism. Superior performance of plants raised from high vigor

**Table 4.** Effect of seed pretreatment with salicylic acid (S.A., 200 µg/ml) and ascorbic acid (A. A., 200 µg/ml) and treatment with *Eucalyptus* oil (E. oil) followed by AA treatment for 20 days on changes of chlorophyll and protein contents in leaves of sunflower plants.

Treatment	Chlorophyll (mg/gfr.wt.)				Protein (mg/gfr.wt.)			
	Plant age (days)							
	30		60		30		60	
	AA (days)							
	0	20	0	20	0	20	0	20
Control	1.20 <sup>a</sup>	0.96 <sup>a</sup>	1.89 <sup>b</sup>	0.88 <sup>a</sup>	15.49 <sup>a</sup>	13.25 <sup>a</sup>	18.25 <sup>a</sup>	16.32 <sup>a</sup>
S.A.	1.43 <sup>b</sup>	1.10 <sup>c</sup>	2.27 <sup>c</sup>	1.83 <sup>d</sup>	21.37 <sup>d</sup>	19.89 <sup>d</sup>	29.60 <sup>d</sup>	26.33 <sup>d</sup>
A.A.	1.26 <sup>a</sup>	1.07 <sup>bc</sup>	2.11 <sup>a</sup>	1.36 <sup>c</sup>	19.22 <sup>c</sup>	17.51 <sup>c</sup>	24.52 <sup>c</sup>	22.63 <sup>c</sup>
E.oil	1.23 <sup>a</sup>	1.00 <sup>a</sup>	1.93 <sup>b</sup>	1.02 <sup>b</sup>	17.33 <sup>b</sup>	15.25 <sup>b</sup>	20.19 <sup>b</sup>	19.16 <sup>b</sup>
LSD(P=0.05)	0.10	0.05	0.13	0.10	1.60	1.33	1.83	1.64

Data were recorded from 30 and 60 days old uniformly grown plants. Values in a vertical column followed by the same letter are not significantly different at 5% level.

**Table 5.** Effect of seed pretreatment with salicylic acid (S.A., 200 µg/ml) and ascorbic acid (A. A., 200 µg/ml) and treatment with *Eucalyptus* oil (E. oil) followed by AA treatment for 20 days on changes of DNA and RNA levels in leaves of sunflower plants.

Treatment	DNA (µg/gfr.wt.)				RNA (µg/gfr.wt.)			
	Plant age (days)							
	30		60		30		60	
	AA (days)							
	0	20	0	20	0	20	0	20
Control	231.1 <sup>b</sup>	182.5 <sup>a</sup>	206.5 <sup>a</sup>	236.1 <sup>b</sup>	905.2 <sup>a</sup>	757.8 <sup>a</sup>	1100.4 <sup>a</sup>	867.0 <sup>cd</sup>
S.A.	238.7 <sup>c</sup>	200.9 <sup>b</sup>	294.8 <sup>d</sup>	244.4 <sup>b</sup>	928.7 <sup>b</sup>	789.4 <sup>c</sup>	1190.5 <sup>d</sup>	948.2 <sup>bc</sup>
A.A.	221.0 <sup>a</sup>	188.9 <sup>a</sup>	265.6 <sup>c</sup>	210.8 <sup>a</sup>	912.4 <sup>a</sup>	783.8 <sup>b</sup>	1168.3 <sup>c</sup>	887.3 <sup>d</sup>
E.oil	217.0 <sup>a</sup>	185.2 <sup>a</sup>	231.7 <sup>b</sup>	207.6 <sup>a</sup>	910.7 <sup>a</sup>	780.5 <sup>b</sup>	1148.7 <sup>b</sup>	815.4 <sup>a</sup>
LSD(P=0.05)	4.01	11.26	18.74	12.88	10.47	4.72	18.54	30.48

Data were recorded from 30 and 60 days old uniformly grown plants. Values in a vertical column followed by the same letter are not significantly different at 5% level.

seeds is available in the literature (Rai, 2000; Pati and Bhattacharjee, 2011). In this investigation, salicylic acid, ascorbic acid and *Eucalyptus* oil-induced enhanced seed germinability, plant growth and metabolism clearly indicate the hardening or invigoration property of the pretreating agents. Such hardening effect on seed was reflected in plant growth and metabolism.

Loss of some vital cellular components occurred (Abdul-Baki and Anderson, 1972; Kole and Gupta, 1982), along with decrease of nucleic acids (Bhattacharjee and Gupta, 1985; Pati, 2007) during the process of seed deterioration are available in literature. Catalase (Abdul-Baki and Anderson, 1972; Yadav et al., 2003) and peroxidase (Bhattacharjee and Choudhuri, 1986; Yadav et al., 2003) activities are generally used as very reliable indices for the evaluation of seed viability. In fact, these two are considered as potential scavenger enzymes which can efficiently detoxify harmful metabolite like H<sub>2</sub>O<sub>2</sub> and thus help alleviating undesired toxic environment at

the cellular level. Hence higher activities of CAT, POD and superoxide dismutase etc. are reported as strong defense elements in plant system (Yadav et al., 2003; Mishra, 2006; Pati, 2007). High level of CAT activity in high vigor seeds have also been reported (Bhattacharjee 1984; Bhattacharjee et al., 1999). *Eucalyptus* oil releasing a number of volatile terpenes and it also contains one active component, eucalyptol which might have a direct role on antimicrobial activity. *Eucalyptus* oil-induced enhancement of seed germinability and metabolism has also been reported and this result is in conformity with reported observation (Singh et al., 1991; Bhattacharjee, 2001; Pati and Bhattacharjee, 2011). So, from the present observations of higher metabolic status of the salicylic acid, ascorbic acid and *Eucalyptus* oil-pretreated sunflower seeds, it seems quite apparent that the seed pretreating agents considerably hardened the seeds and such hardening is effected at the metabolic level which subsequently resulted in retention of seed

**Table 6.** Effect of seed pretreatment with salicylic acid (S.A., 200 µg/ml) and ascorbic acid (A. A., 200 µg/ml) and treatment with *Eucalyptus* oil (E. oil) followed by AA treatment for 20 days on changes of catalase and peroxidase activities in leaves of sunflower plants.

Treatment	Catalase (unit/h/gfr.wt.)				Peroxidase (unit/h/gfr.wt.)			
	Plant age (days)							
	30		60		30		60	
	AA (days)							
	0	20	0	20	0	20	0	20
Control	100.4 <sup>a</sup>	68.9 <sup>a</sup>	163.6 <sup>a</sup>	102.7 <sup>a</sup>	94.8 <sup>a</sup>	59.7 <sup>a</sup>	145.9 <sup>a</sup>	97.7 <sup>a</sup>
S.A.	112.3 <sup>b</sup>	82.0 <sup>c</sup>	184.2 <sup>b</sup>	122.2 <sup>c</sup>	109.2 <sup>b</sup>	74.6 <sup>c</sup>	168.7 <sup>b</sup>	135.4 <sup>c</sup>
A.A.	111.4 <sup>b</sup>	75.3 <sup>b</sup>	166.0 <sup>a</sup>	110.3 <sup>b</sup>	101.1 <sup>a</sup>	68.5 <sup>b</sup>	154.6 <sup>a</sup>	126.0 <sup>b</sup>
E.oil	108.7 <sup>b</sup>	69.8 <sup>a</sup>	162.3 <sup>a</sup>	103.5 <sup>a</sup>	98.7 <sup>a</sup>	63.1 <sup>ab</sup>	151.1 <sup>a</sup>	121.6 <sup>b</sup>
LSD(P=0.05)	6.05	5.10	12.40	6.33	7.08	6.07	10.05	7.09

Data were recorded from 30 and 60 days old uniformly grown plants. Values in a vertical column followed by the same letter are not significantly different at 5% level.

**Table 7.** Effect of seed pretreatment with salicylic acid (S.A., 200 µg/ml) and ascorbic acid (A. A., 200 µg/ml) and treatment with *Eucalyptus* oil (E. oil) followed by AA treatment on yield attributes of sunflower plants.

Treatment	Capitulum diameter (cm)		Capitulum fr.wt.(g)		Seed no./capitulum		1000Seedwt. (g)	
	AA (days)							
	0	20	0	20	0	20	0	20
Control	14.0 <sup>a</sup>	10.3 <sup>a</sup>	212.6 <sup>a</sup>	164.9 <sup>a</sup>	1001.5 <sup>a</sup>	764.1 <sup>b</sup>	38.4 <sup>a</sup>	32.2 <sup>a</sup>
S.A.	15.9 <sup>b</sup>	14.1 <sup>c</sup>	249.0 <sup>b</sup>	181.2 <sup>b</sup>	1103.0 <sup>b</sup>	821.0 <sup>a</sup>	48.0 <sup>c</sup>	43.0 <sup>a</sup>
A.A.	14.8 <sup>a</sup>	13.0 <sup>d</sup>	220.1 <sup>a</sup>	177.0 <sup>b</sup>	1012.0 <sup>a</sup>	816.7 <sup>a</sup>	46.2 <sup>c</sup>	38.1 <sup>c</sup>
E.oil	14.0 <sup>a</sup>	11.2 <sup>a</sup>	215.3 <sup>a</sup>	160.6 <sup>a</sup>	1006.9 <sup>a</sup>	800.5 <sup>a</sup>	42.7 <sup>b</sup>	35.6 <sup>b</sup>
LSD(P=0.05)	1.03	1.01	15.33	12.36	68.51	65.20	2.63	2.10

Data were recorded after harvest. Values in a vertical column followed by the same letter are not significantly different at 5% level.

vigor and consequent extension of seed viability with concomitant enhancement of plant potential.

From the overall results of this study, it can be concluded that salicylic acid, ascorbic acid and *Eucalyptus* oil are not only effective in enhancing storage potential of sunflower seeds but they are equally efficient in improving plant vigor with concomitant augmentation of crop yield as evident from the results which clearly showed that the chemical-treated seeds were less sensitive than the control ones under adverse storage situations. What ever might be the mechanism for the chemical manipulated seed invigoration, results of this investigation can at least indicate that the experimental chemicals has some efficacy for enhancement of storage potential of sunflower seeds and field performance of plants raised from such potentiated seeds of the test species. The results also pointed out that the catabolic process in treated seeds remained partially subdued, there by rendering the seeds against unfavorable storage environment. And thus, the invigoration property of the present seed pretreating agents seems to be apparent from the experimental results.

## REFERENCES

- Abdul-Baki AA, Anderson JD (1972). Physiological and biochemical deterioration of seeds. In: Kozlowski T (ed.), Seed Biology, 2: 203-215.
- Anderson JD (1970). Metabolic changes in partially dormant wheat seeds during storage. Plant Physiol., 46: 605-608.
- Arnon DI (1949). Copper enzymes in isolated chloroplasts. Polyphenol oxidase in *Beta vulgaris*. Plant Physiol., 24: 1-15.
- Aziz NH, Shair AAM (1979). Influence of other fungi on aflatoxin production by *Aspergillus flavus* in maize kernels. J. Food Safety 17(2): 113-123.
- Basu RN (1994). An appraisal of research on wet and dry physiological seed treatments and their applicability with special reference to tropical and subtropical countries. Seed Sci. Technol., 22: 107-1261.
- Bhattacharjee A (1984). Responses of sunflower plants towards growth retardants with special reference to growth, metabolism and yield. Ph.D. Thesis, Burdwan University, India.
- Bhattacharjee A, Choudhuri MA (1986). Chemical manipulation of seed longevity and stress tolerance capacity of seedlings of *Corchorus capsularis* and *C. olitorius*. J. Plant Physiol., 125: 391-400.
- Bhattacharjee A, Gupta K (1985). Effect of dikegulac sodium, a growth retardant, on the viability of sunflower seeds. Seed Sci. Technol., 13: 165-174.
- Bhattacharjee A, Chattopadhyay A, Rai AS (1999). Potentiation of safflower seeds under artificial stress storage environment by chemical growth retardants. Seed Sci. Technol., 27: 707-719.

- Bhattacharyya AK, Basu RN (1990). Retention of vigour and viability of stored pea (*Pisum sativum* L.) seed. *Indian Agriculturist*, 34: 187-193.
- Biswas AK, Choudhuri MA (1978). Differential behaviour of the flag leaf of intact rice plant during ageing. *Biochem. Physiol. Pflanzen*, 173: 220-228.
- Bhattacharjee A (2001). Influence of growth retardant and a volatile oil and enhancement of storage longevity of two low vigour seed species. *J. National Botanic. Soc.*, 55: 55-61.
- Barman P (2006). Seed parameters study, prediction and enhancement of seed yield in different genotypes of *Vigna* species. Ph.D. Thesis, Bidhan Chandra Krishi Vishwavidyala, West Bengal, India.
- Cherry JH (1962). Nucleic acid determination in storage tissue of higher plants. *Plant Physiol.*, 37: 670-678.
- Chhetri DR, Rai AS, Bhattacharjee A (1993). Chemical manipulation of seed longevity of four crop species in an unfavourable storage environment. *Seed Sci. Technol.*, 21: 31-44.
- Choudhuri MA, Chatterjee SK (1970). Seasonal changes in the level of some cellular components in the abscission zone of *Coleus* leaves of different ages. *Annal. Bot.*, 34: 275-287.
- Christensen CM, Kaufmann HH (1965). Deterioration of stored grain by fungi. *Annal Rev. Phytopathol.*, 3 : 69-84.
- Copland LO, McDonald MB (2001). In *Principles of Seed Sc. and Technol.* (4th edn.), Norwell, Massachusetts: Kluwer Academic Publishers, USA.
- Delouche JC, Baskin CC (1973). Accelerated ageing techniques for predicting the relative storability of seed lots. *Seed Sci. Technol.*, 1: 427 – 452.
- Fick NG, Qualset CO (1975). Genetic control of endosperm amylase activity and gibberellin responses in standard height and short statured wheat. *Proceedings of National Academy of Science. USA.* 72: 892-895.
- Halder S (1981). Studies on viability, yield and associated biochemical changes in leaves during seed filling in sunflower (*Helianthus annuus* L. cv. EC 68414). Ph.D. Thesis, Burdwan University, India.
- Halder S, Koley S, Gupta K (1983). On the mechanism of sunflower seed deterioration under two different types of accelerated ageing. *Seed Sci. Technol.*, 11: 331-339.
- Heydecker W (1972). Vigour. In: E.H. Roberts (ed.), *Viability of Seeds*, Chapman and Hall Ltd., London, pp. 209-252.
- International Seed Testing Association, ISTA (1976). International rules for seed testing. *Seed Sci. Technol.*, 4: 51-177.
- Kar M, Mishra D (1976). Catalase, peroxidase, polyphenol oxidase activities during rice leaf senescence. *Plant Physiol.*, 57: 315-319.
- Kole S, Gupta K (1982). Biochemical changes in safflower (*Carthamus tinctorius*) seeds under accelerated ageing. *Seed Sci. Technol.*, 10: 47-54.
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ (1951). Protein measurement with the folin-phenol reagent. *J. Biol. Chem.*, 193: 265-275.
- Markham R (1955). Nucleic acids, their compounds and related compounds. In: K. Peach and M.V. Tracey (eds.), *Modern Methods of Plant Analysis*, Springer Verlag, Berlin, 4: 246-304.
- Mishra VK (2006). Studies on seed vigour and viability of a few pulse crops and devising techniques of seed invigouration. Ph.D. Thesis, Vidyasagar University, India.
- Panase VG Sukhatme PT (1967). *Statistical Methods for Agricultural Workers*. 2nd edition, ICAR, New Delhi, pp. 150-157.
- Perl M, Luria I, Gelmond H (1978). Biochemical changes in sorghum seeds affected by accelerated ageing. *J. Expt. Bot.* 29: 497-509.
- Pati CK (2007). Seed invigouration, plant potentiation and yield augmentation of two promising pulse crops (*Lathurus sativus* L. and *Vinna mungo* (L.) Hepper) by chemical manipulation. Ph.D. thesis, Vidyasagar University, West Bengal, India.
- Pati CK, Bhattacharjee A (2011). Alleviation of seed deterioration and enhancement of plant potential of a pea species using selected medicinal plant extracts. *J. Med. Plant Res.*, 5(17): 4108-4111.
- Rai A (2000). *An investigation into the problems of maintenance of seed vigour and viability under adverse climatic conditions of Darjeeling hills*. Ph.D. Thesis, North Bengal University, India.
- Rao RGS, Singh PM, Singh J, Singh KP (2003). Deterioration of onion seeds under accelerated ageing conditions. *Indian J. Plant Physiol. Special Issue*: 189-194.
- Roosrokh M, Golozani KG, Javanshir A (2002). Relationship between seed vigour and field performance in chick pea (*Cicer arietinum* L.). *Seed Plant*, 18: 156-169.
- Snell FD, Snell CT (1971). *Colorimetric Methods of Analysis*. Van Nostrand Reinhold Co., New York, IV AAA, pp. 7-145.
- Sengupta AK, De BK, Mondal AK (2005). Pre-storage seed invigouration treatments for the maintenance of vigour, viability and field performance of high-vigour onion seed (*Allium cepa* L.). *Seed Sc. Technol.*, 33: 753-760.
- Singh S, Kohli RK, Saxena DB (1991). Effect of eucalyptus oil on germination and growth of *Phaseolus aureus* Roxb. *Plant Soil.*, 137(2): 223-227.
- Yadav S, Bhatia VS, Guruprasad KN (2003). Role of peroxidase and catalase enzymes in deterioration of soyabean seeds due to field weathering. *Indian J. Plant Physiol., Special Issue*, pp. 195-200.