

*Full Length Research Paper*

# Knowledge and awareness on HIV/AIDS among blood donors: A Study at Rajshahi, Bangladesh

Abdul Goni\* and Mahfuzur Rahman

Department of Population Science and HRD, University of Rajshahi, Rajshahi-6205, Bangladesh.

Accepted 17 November, 2017

Human immunodeficiency virus/acquired immune deficiency syndrome (HIV/AIDS) is the most devastating disease that mankind has ever faced. For being neighboring country of India, Bangladesh has been highly vulnerable to HIV infection. Though Bangladesh continues to maintain low HIV prevalence status, however the infection rate is on the rise at Rajshahi Division. Some studies have been carried out on the knowledge about HIV/AIDS among some risk groups, but studies on blood donor's awareness about HIV/AIDS are rarely found. This study aims at investigating the knowledge and awareness about HIV/AIDS among the blood donors at Rajshahi Metropolitan of Bangladesh. In this study, primary data was used. Data reveals that, though about 89% blood donors knew it is transmitted through blood transfusion, only 1% of them had undertaken blood test before donation, which is alarming. The odds of having preliminary and preventive knowledge fell with the increase in age of the respondents. Besides these, blood donors who were educated and who were in service were found more aware about the HIV/AIDS than the others. Strict rules and regulations should be maintained regarding blood screening. Overall people should be encouraged to use condoms and abide by the religious rules to avoid HIV/AIDS infection. Both, Government and Non-Government organizations should augment their educational and focused group discussion (FGD) programs on HIV/AIDS knowledge.

**Key words:** Human immunodeficiency virus (HIV), acquired immune deficiency syndrome (AIDS), blood donors, awareness, focused group discussion (FGD).

## INTRODUCTION

AIDS stands for acquired immune deficiency syndrome. The human immunodeficiency virus, which is commonly called HIV, is a virus that directly attacks certain human organs, such as the brain, heart, and kidneys, as well as the human immune system. The immune system is made up of special cells, which are involved in protecting the body from infections and some cancer. HIV damages the natural defense system in human body and the state is called AIDS.

HIV/AIDS is the most devastating disease that mankind has ever faced. HIV does not survive well outside the

body. It can not be transmitted through casual daily contact. Mosquitoes and other insects do not transmit HIV. HIV can be spread by sexual contact with an infected person, by sharing needles and/or syringes and/or other injecting equipments or, less commonly (and now very rarely in countries where blood is screened for HIV antibodies), through transfusion of infected blood or blood clotting factors. Babies born to HIV infected women may become infected before or during birth or through breastfeeding after birth.

After the first detection of AIDS in 1981 in New York City and San Francisco, this disease in Bangladesh was first found in 1983. Since the first detection of HIV in Bangladesh, the rate of infection has not increased in comparison to our neighbors (for example, India). Given that the overall prevalence of HIV remains below 1%

\*Corresponding author. E-mail: [magoni\\_popsrbd@yahoo.com](mailto:magoni_popsrbd@yahoo.com).  
Tel: +88 0721 750041/4121, +88.

(0.7%). Bangladesh continues to maintain low prevalence status (National HIV serological surveillance, Bangladesh, 9<sup>th</sup> round, 2011). But it is clear that this situation may not continue if the risky behaviors that increase vulnerability cannot be contracted among the high-risk group, vulnerable group and general population. Needle sharing among drug users occurs at alarmingly high rates (77%) in central region (National HIV serological surveillance, Bangladesh, 4<sup>th</sup> round, 2002). of Bangladesh. In a study, Rahman et al. (1998) revealed that only 18% Commercial Sex Workers (CSW) in Bangladesh heard about AIDS and 8% knew that it can be transmit by sexual contact. AIDS was believed to be curable by 5% of the total respondents. A vigorous AIDS awareness campaigns for the CSWs and their clients is necessary to prevent an explosive epidemic in Bangladesh. In another study, Kumar et al. (2002) concluded that awareness about HIV/AIDS infection, transmission and preventive measures is the essential thing for the people in the countries of developing world, especially for women who are engaged in commercial sex works and those of socially and deprived groups. Indeed, there is a host of factors that have rendered the country highly vulnerable to a surge in the epidemic. These include the overall poverty levels, the documented risk behaviors including needle sharing, growing sex work, population mobility, persisting gender disparity and inequality, low levels of general awareness and knowledge about HIV/AIDS among general population, and unscreened blood transfusion.

In many countries, it has been observed that, once an HIV epidemic begins among Injection Drug Users (IDU), it is unlikely to remain limited in that population (Cohen, 2004; Panda, 2000). This is especially true in Bangladesh, since behavior surveillance data show that IDU are not at all an isolated population. Drug injector's sells blood and highly involved in other risk behaviors. HIV prevalence among IDUs has increased to 5.3% in 2011 (National HIV serological surveillance, Bangladesh, 9<sup>th</sup> round, 2011). From 4% in 2002 (National HIV serological surveillance, Bangladesh, 4<sup>th</sup> round, 2002) which is a threat for blood receivers, as blood screening system is very poor in Bangladesh. According to the Federal Centers for Disease Control and Prevention (CDC), about 6% infection among women occurs through blood transfusion, which is threatening to the future generation. Bangladesh professional donor should have knowledge about HIV/AIDS. Since Bangladesh is a developing country, most of the people do not have enough knowledge about the importance of blood test before blood donation. A limited number of blood donors have knowledge about HIV/AIDS in Bangladesh.

Though it is encouraging that, HIV prevalence has declined over the years among population at risk in some geographic areas of Bangladesh, however in many areas

especially in Rajshahi Division, high and rising HIV prevalence was recorded (National HIV serological surveillance, Bangladesh, 9<sup>th</sup> round, 2011). The rise of infection at Rajshahi has put the city into the category of vulnerable area. Some researchers (Rahman et al., 1998; Khan et al., 1997) have studied the knowledge about HIV/AIDS among some risk groups, such as patients with sexually transmitted disease, pregnant woman at antenatal clinics, commercial sex workers, long distant truck driver and overseas workers. But the studies on awareness about HIV/AIDS of blood donors are very rare. The purpose of this study is to focus the knowledge and awareness scenario on different aspects of HIV/AIDS among the blood donors at Rajshahi Metropolitan (divisional city) of Bangladesh.

## MATERIALS AND METHODS

This study is based on primary data. The data was collected under the authority of the Department of Population Science and Human Resource Development of Rajshahi University, Bangladesh. To collect the data, purposive sampling technique was applied. Interview method has been used to collect information from the respondents. A total of 160 blood donors of Rajshahi Metropolitan had been interviewed and their information was recorded. The period of data collection was 3rd March to 28 April in 2008. Among the total respondents, only 21% were female and 52% were married. Only 18% blood donors were under 21 years of age, while 4% were above age 40 years. 88% donors were Muslim and 64% were village dweller. It was observed that, 49% blood donors did not have any income, as most of them were student and only 2% had income over 20000 Tk/month (about US \$ 250). 12% donors were illiterate, while 7% were with post graduation. Among the respondents, 34% were student, 16% were agriculture worker, and more interestingly 13% were house wife.

In this study, descriptive statistical techniques have been applied to examine the differentials. Besides the descriptive statistics, multivariate analytical technique (logistic regression model) has also been used to investigate the determinants of knowledge and awareness about HIV/AIDS. The statistical analyses have been performed by the software SPSS. In this study, two sets of variables are selected as dependent variable, which are variables relating preliminary knowledge of HIV/AIDS (which includes whether knows the name of HIV/AIDS, agree that AIDS is a fatal disease, AIDS is not a contagious disease and knowledge about the medium of HIV/AIDS infection), and variables relating preventive knowledge of HIV/AIDS (includes the knowledge about the means of preventing HIV/AIDS infection). The variables relating preliminary knowledge and preventive knowledge were further categorized into two groups depending upon the extent of knowledge, the categories are sufficient and insufficient knowledge. The classification was made by subjective judgment. For both the variables, if the 90% of the answers of the respondent were found correct then the respondent was considered having sufficient knowledge, otherwise he was considered to have insufficient knowledge. Independent variables used in the analyses are age, sex, education, occupation, possession of Radio/TV, condom's impact and religious rule's impact. In case of independent variable, classifications for bivariate analysis differ from that of multivariate analysis, where in multivariate analysis; independent variables include lower number of classes than that in bivariate analysis. This is done to make the

variation of impact of different classes more clear and apparent, and to facilitate interpretability.

In multivariate analysis, logistic regression model (Cox, 1972) has been applied. This model can be used to identify the risk factors as well as to predict the probability of success e.g. probability of developing a disease as a function of the particular risk factor. This probability can serve as an index of risk for a given disease or for not responding to certain treatment. The logistic regression has become the standard method for finding the relationship between the qualitative outcome variables and a set of explanatory variables. Then the logit transformation is defined to be

$$z_i = \log \text{it} (p_i) = \left( \frac{p_i}{1-p_i} \right) = x_i \beta$$

Where  $\beta$ , a regression coefficient and this equation are known as the logit model that relates the independent variables to the transformation of  $p_i$ .

Taking these probabilities ( $p$  and  $1-p$ ) as the basis of analysis, some functions are considered that transform the scale (0, 1) for the probabilities on to the real line. This function is known as link functions or response function. The logit model can be written as

$$\log \text{it} (p_i) = \log [p_i / (1 - p_i)] = X_i \beta = \beta_0 + \beta_1 X_{1i} + \beta_2 X_{2i} + \dots + \beta_k X_{ki}$$

Where  $p_i = \Pr(Y_i = 1 | x_i) = \frac{\exp(X_i \beta)}{1 + \exp(X_i \beta)}$

## RESULTS AND DISCUSSION

To study the blood donor's knowledge and awareness about HIV/AIDS, both, bivariate and multivariate analytical techniques have been used. To dissect the knowledge and awareness scenario and their pattern, descriptive statistical technique has been applied, and to spot the significant covariates and their degrees of impact on the knowledge and awareness of blood donors about HIV/AIDS, Logistic Regression analysis has been used. The results and discussion have been presented in the following sections.

### Knowledge and awareness about HIV/AIDS

Main objective of this paper is to investigate the blood donor's knowledge and awareness about HIV/AIDS. Several questions were tried to capture their knowledge and awareness which have been summarized in Table 1.

From the table, it is observed that 97% blood donors heard the name of AIDS and 76% knew it is caused by HIV virus, whereas 3 and 24% respectively never heard the name of AIDS and do not know that it is caused by HIV virus. 94% blood donors knew it is fatal and 89% knew that it is transmitted through blood transfusion. But the worst thing is that, 94% blood donors did not test their blood before donating blood; only 1% did the test.

89% knew that HIV/AIDS transmits through sexual intercourse with more than one partner, 87% knew it is transmitted through sharing needle with others, 66% knew it may spread from pregnant mother to her child and 52% knew it may transmit through breast feeding.

Regarding preventive knowledge, 8% knew that it may be prevented by vaccine, 89% knew it can be prevented by using condom during sexual intercourse. 93% believed by increasing mass awareness and 92% believed by abiding religious rules, HIV/AIDS can be prevented.

### Level and pattern of awareness and knowledge of the blood donors by background characteristics

Table 3 shows that, 67 respondents (41%) were found to have sufficient preliminary knowledge about HIV/AIDS and 109 respondents (68%) having sufficient knowledge about preventive measures of HIV/AIDS. Respondents of age group 21 to 25 years were most aware about both preliminary (22 out of 67; 33%) and preventive (35 out of 109; 32%) knowledge about HIV/AIDS. It was observed that the awareness decreases with the increase in age. Male respondents were found more aware about preliminary (57 out of 67; 85%) and preventive knowledge (89 out of 109; 81%) than its female counterpart. A positive relationship has been observed between educational level and awareness as in Table 2.

It was observed that, those who are student and service holder are more aware about HIV/AIDS. Those having Radio/Television were more conscious about the preliminary and preventive knowledge about HIV/AIDS. Higher percentage is observed to have sufficient knowledge about HIV/AIDS for those who know that using condom and abiding by religious rules can prevent HIV/AIDS than those who do not know.

Given the descriptive statistics and information of the different explanatory variables above, which covariates are likely to affect knowledge and awareness of blood donors comes into question? To identify the influential factors, Logistic regression analysis was performed and the result has been presented in the next section.

### Logistic regression analysis

The logistic regression analysis is aimed at identifying the important contribution of variables that have an influence on the preliminary and preventive knowledge of blood donors. A variable was considered significantly associated with having knowledge, if its P value was below 0.10. The results of the analyses are shown in Table 3. From the analysis, we observed that, respondents of age group 20 to 24 years are 2.2 and 1.2

**Table 1.** Percentage distribution of respondents by selected categories having knowledge and awareness about HIV/AIDS.

<b>Questions</b>	<b>No.</b>	<b>Age (%)</b>
<b>Did you listen the name HIV/ AIDS?</b>		
Yes	153	95.6
No	7	4.4
<b>Do you know the name of virus which causes AIDS?</b>		
Yes	122	76.3
No	31	19.4
No answer	7	4.4
<b>Is AIDS a fatal disease?</b>		
Yes	151	94.4
No	2	1.3
No answer	7	4.4
<b>Does HIV transmit through blood transfusion?</b>		
Yes	142	88.8
No	3	1.9
Doesn't know	8	5.0
No answer	7	4.4
<b>Before donation did you test blood?</b>		
Yes	2	1.3
No	151	94.4
No answer	7	4.4
<b>Does HIV/AIDS transmit through sexual intercourse with more than one male or female?</b>		
Yes	143	89.4
Doesn't know	10	6.3
No answer	7	4.4
<b>Does HIV/AIDS transmit through sharing same needle with others?</b>		
Yes	139	86.9
No	1	0.6
Doesn't know	13	8.1
No answer	7	4.4
<b>Does HIV/AIDS transmit from pregnant mother to her child?</b>		
Yes	107	66.9
No	3	1.9
Doesn't know	43	26.9
No answer	7	4.4
<b>Does HIV/AIDS transmit through breast feeding?</b>		
Yes	83	51.9
No	4	2.5
Doesn't know	66	41.3
No answer	7	4.4

Table 1. Contd.

<b>Have any vaccine to prevent HIV/AIDS?</b>			
Yes		12	7.5
No		74	46.3
Doesn't know		66	41.3
No answer		8	5.0
<b>Do you know that HIV/AIDS can be prevented by using condom?</b>			
Yes		143	89.4
Doesn't know		10	6.3
No answer		7	4.4
<b>Do you think that HIV/AIDS can be prevented by mass awareness?</b>			
Yes		148	92.5
Doesn't know		5	3.1
No answer		7	4.4
<b>Do you think that HIV/AIDS can be prevented by abiding religious rules?</b>			
Yes		147	91.9
Doesn't know		6	3.8
No answer		7	4.4

Rajshahi field data, April, 2008.

Table 2. Percentage distribution regarding sufficient and insufficient preliminary knowledge and preventive knowledge about HIV/AIDS of the respondents.

Characteristics	Preliminary knowledge about HIV/AIDS		Preventive knowledge about HIV/AIDS	
	Sufficient knowledge (%)	Insufficient knowledge (%)	Sufficient knowledge (%)	Insufficient knowledge (%)
<b>Age (in years)</b>				
<21	16 (55.2)	13 (44.8)	23 (79.3)	6 (20.7)
21-25	22 (38.6)	35 (61.4)	35 (61.4)	22 (38.6)
26-30	13 (40.6)	19 (59.4)	23 (71.9)	9 (28.1)
31-35	10 (55.6)	8 (44.4)	14 (77.8)	4 (22.2)
36-40	5 (29.4)	12 (70.6)	11 (64.7)	6 (35.3)
40 <sup>+</sup>	1 (14.3)	6 (85.7)	3 (42.9)	4 (57.1)
<b>Sex</b>				
Male	57 (44.9)	70 (55.1)	89 (70.1)	38 (29.9)
Female	10 (30.3)	23 (69.7)	20 (60.6)	13 (39.4)
<b>Education</b>				
Illiterate	2 (10.5)	17 (89.5)	7 (36.8)	12 (63.2)
Primary	7 (23.3)	23 (76.7)	13 (43.3)	17 (56.7)
Secondary	9 (31.0)	20 (69.0)	17 (58.6)	12 (41.4)
Higher Secondary	17 (51.5)	16 (48.5)	28 (84.8)	5 (15.2)
Graduate	24 (63.2)	14 (36.8)	33 (86.8)	5 (13.2)
Post-Graduate	8 (72.7)	3 (27.3)	11 (10)	0 (0)
<b>Occupation</b>				
Student	33 (60.0)	22 (40.0)	46 (83.6)	9 (16.4)

Table 2. Contd.

Agriculture	3 (12.0)	22 (88.0)	10 (40.0)	15 (60.0)
Service	20 (69.0)	9 (31.0)	25 (86.2)	4 (13.8)
Business	7 (23.3)	23 (76.7)	18 (60.0)	12 (40.0)
Housewife	4 (19.0)	17 (81.0)	10 (47.6)	11 (52.4)
<b>Having Radio/TV</b>				
Yes	40 (39.2)	31 (53.4)	61 (59.8)	10 (17.2)
No	27 (46.6)	62 (60.8)	48 (82.8)	41 (40.2)
<b>HIV/AIDS can be prevented by using condom</b>				
Yes	57 (40.1)	85 (59.9)	95 (69.3)	42 (30.7)
No	10 (55.6)	8 (44.4)	14 (60.9)	9 (39.1)
<b>HIV/AIDS can be prevented by abiding religious rules</b>				
Yes	61 (43.6)	79 (56.4)	99 (71.2)	40 (28.8)
No	6 (30)	14 (70)	10 (47.6)	11 (52.4)
Total	67 (41.9)	93(58.1)	109(68.1)	51(31.9)

Rajshahi field data, April, 2008.

times more likely to have preliminary and preventive knowledge about HIV/AIDS respectively than the respondent of below 20 years of age. But the odds fell with the increase in age. In the analysis, male respondent received 2.5 and 1.3 times higher odds than its female counterpart to have preliminary knowledge and preventive knowledge respectively about HIV/AIDS. Respondents with primary and secondary education and higher education received higher odds than the illiterate respondents. Service holders were 100 and 83% more likely to have preliminary and preventive knowledge respectively about HIV/AIDS than the others. Respondents had radio/TV were more likely to have knowledge about HIV/AIDS than those did not have any radio/TV. Persons knew that HIV/AIDS can be prevented by using condoms were 5.17 and 2.91 times more likely to possess preliminary and preventive knowledge about HIV/AIDS. Among those who knew that HIV/AIDS can be prevented by abiding by religious rules received higher odds of having preliminary and preventive knowledge than those who did not know.

## Conclusion

HIV/AIDS is one of the deadliest disease that does not have any treatment, thus to get rid of the catastrophe, the only way out there is its prevention. To prevent the spread of the disease, knowledge and awareness among

the high risk group about it is indispensable. This study attempts to examine the knowledge and awareness of blood donors about HIV/AIDS. The matter of hope is that, 89% blood donors know that HIV/AIDS is transmitted by blood transfusion, but the worst thing is that only 1% undertook blood test before donation, which is frustrating. A sizeable magnitude knows that it can be prevented by using condom (89%) and by abiding religious rules (92%). In this study, age, sex, education, occupation, Radio/TV possession and knowledge about preventive role of condom and religious rules have been deemed to be important determinants of knowledge about both preliminary and prevention of HIV/AIDS. Interestingly, the awareness decreases with the increase of age, while all other variables are found positively related with awareness. To increase awareness in a particular population segment, focused group discussion (FGD) is the most effective tool and considerable emphasize should be given to FGD. Undoubtedly education is the most important determinant of any sort of awareness, and so in this study. GOs and NGOs should augment their educational program containing knowledge about HIV/AIDS, throughout the country.

Finally, radio and TV are the most popular medium of recreation, programs relating preliminary and prevention knowledge about HIV/AIDS should be increased in radio and TV. Besides these, overall People should be encouraged to use condoms and abide by the religious rules by the popular media. Now under the HAPP 98 safe

**Table 3.** Multivariate logistic regression estimates of relative odds of having preliminary knowledge and preventive knowledge about HIV/AIDS.

Characteristics	Relative risk (odd ratio)	
	Having preliminary knowledge about HIV/AIDS	Having preventive knowledge about HIV/AIDS
<b>Age (in years)</b>		
<20 (RC)	1.00 <sup>***</sup>	1.00 <sup>***</sup>
20-24	2.22 <sup>^^</sup>	1.23 <sup>^^</sup>
25-29	1.06 <sup>^^</sup>	1.39 <sup>^^</sup>
30-34	1.82 <sup>***</sup>	2.19 <sup>***</sup>
35 <sup>+</sup>	1.23 <sup>^</sup>	2.05 <sup>^</sup>
<b>Sex</b>		
Female (RC)	1.00	1.00
Male	2.51 <sup>**</sup>	1.31 <sup>**</sup>
<b>Education</b>		
Illiterate (RC)	1.00 <sup>***</sup>	1.00 <sup>***</sup>
Primary and Secondary	1.30 <sup>***</sup>	1.68 <sup>***</sup>
Higher	1.86 <sup>^</sup>	2.29 <sup>^</sup>
<b>Occupation</b>		
Others (RC)	1.00 <sup>***</sup>	1.00 <sup>***</sup>
Service Holder	2.03 <sup>^</sup>	1.83 <sup>^</sup>
<b>Having Radio/TV</b>		
No (RC)	1.00 <sup>***</sup>	1.00 <sup>***</sup>
Yes	2.63 <sup>^</sup>	1.72 <sup>^</sup>
<b>Condom can prevent HIV/AIDS</b>		
Doesn't Know (RC)	1.00	1.00
Know	5.17 <sup>**</sup>	2.91 <sup>**</sup>
<b>Abiding religious rules can prevent HIV/AIDS</b>		
Doesn't Know (RC)	1.00 <sup>^</sup>	1.00 <sup>^</sup>
Know	2.84 <sup>^</sup>	3.20 <sup>^</sup>

Rajshahi field data, April, 2008; RC=Reference Category, \* = P<0.10, \*\* = P<0.05, \*\*\* = P<0.01.

blood transfusion centers are screening the blood for HIV and 5 other diseases. Government of the country should pay special attention to the blood screening system. Strict rule should be established to screen the blood before taking that from the donors. The blood screening service should be expanded to Thana (sub-district) level and the numbers of the screening centers should be increased in metropolitan cities like Rajshahi.

As HIV/AIDS is an incurable disease, massive steps should be taken to prevent it through building awareness among the blood donors. Some recommendations regarding increase in knowledge and awareness level have been pointed out above which could be considered by the policy makers to improve the scenario.

## REFERENCES

- Cohen J (2004). Asia and Africa: On Different Trajectories. *Science*, 304: 1932-1938.
- Cox DR (1972). Regression and Life Tables. *J. Royal Stat. Soc.*, 34(B): 187-220.
- Khan MA, Rahman M, Khanam PA, Barkat-e-khuda, Kano TT, Ashraf A (1997). Awareness of Sexually Transmitted Diseases among Women and Service Providers in Rural Bangladesh. *INTJ STD AIDS*, 8: 688-690.
- Kumar MN, Shiva R, Raju KNM (2002). Awareness knowledge and misconception about HIV/AIDS: A comparison of two states in South India. *Popul. Stabilization Dev.*, pp. 720-748.
- National HIV serological and Behaviour surveillance (2002). Fourth round technical report. National AIDS/STD Programme, Ministry of Health and Family Welfare, Govt. of the Peoples Republic of Bangladesh.

National HIV serological surveillance, Bangladesh (2011). Ninth round technical report; IEDCR and icddr, b. National AIDS/STD Programme, Directorate General of Health Services, Ministry of Health and Family Welfare, Govt. of the Peoples Republic of Bangladesh.

Panda S (2000). Transmission of HIV from injecting drug users to their wives in India. *Int. J. STD AIDS*, 11: 468-473.

Rahman M, Wali-UI-Islam M and Fukua T. Knowledge and Practice about HIV/AIDS among the Commercial Sex Workers in Bangladesh. *J. Epidemic*. 1998; 8: 256.