

Full Length Research Paper

An investigation on the dynamics of insecticide resistance in *Anopheles gambiae* from southern Benin and the metabolic resistance mechanisms involved

*Gildas Armel, Modeste Fabrice and Hervé Carlos

Department of Parasitology/Microbiology, National University of Benin, Abomey Calavi, Benin.

*Corresponding author. E-mail: gildas.armel@yahoo.com

Accepted 17 December, 2014

Permanet 2.0 distribution was made free in October, 2008 and May, 2009 in Oueme Department. OlysetNet distribution was also made free in July, 2011 throughout the entire country by Beninese National Malaria Control Programme to increase coverage of long-lasting insecticidal nets (LLINs). We investigated the dynamics of insecticide resistance in *Anopheles gambiae* from southern Benin and the metabolic resistance mechanisms involved. Larvae and pupae of *A. gambiae* s. l. were collected from the breeding sites in Littoral and Oueme Departments. Centers for Disease Control and Prevention (CDC) susceptibility tests were conducted on unfed female mosquitoes aged 2 to 5 days old with stock solutions of permethrin, deltamethrin and dichlorodiphenyltrichloroethane (DDT). CDC biochemical assays using synergists were also carried out. *A. gambiae* Ladj populations were resistant to permethrin and DDT in 2008 and in 2013. *A. gambiae* Sekandji populations were susceptible to deltamethrin in 2008 and in 2013 whereas these populations were resistant to this product in 2010. *A. gambiae* Sekandji populations were resistant to DDT in 2008 and in 2013. The DDT resistance level in *A. gambiae* Ladj and Sekandji populations recorded in 2013 was higher than the one observed in 2008. The metabolic resistance conferred by detoxifying enzymes is an indication of phenotypic resistance to both DDT and pyrethroids.

Key words: Dynamics, piperonyl butoxide, S.S.S-tributylphosphorotrithioate, ethacrynic acid, insecticide, vectors, resistance.

INTRODUCTION

Malaria vector control in Africa relies heavily on the organochlorine, dichlorodiphenyltrichloroethane (DDT) and

pyrethroid insecticides such as permethrin, deltamethrin (Zaim et al., 2000). The first cases of pyrethroid resistance

were recorded in West Africa more precisely in Côte d'Ivoire (Elissa et al., 1993). Many other cases have been described in West Africa (Akogbeto and Yakoubou, 1999), in Benin (Chandre et al., 1999), in Burkina Faso (Chandre et al., 1999; Diabaté, 1999) and in Mali (Fanello et al., 2003). In West Africa, the first cases of dieldrin resistance in *Anopheles gambiae* were recorded in Burkina Faso in 1960 (Coz et al., 1968). Ten years later, the identification of cases of DDT resistance was reported in Togo, Senegal, Nigeria (OMS, 1976). In Benin, Akogbeto and Yakoubou (1999) suspected the emergence of DDT resistance recorded in *A. gambiae* from meridian regions, to be related to two phenomena: (i) the massive use of DDT and dieldrin for house-spraying applications in southern villages from 1953 to 1960 during World Health Organization (WHO) programmes of malaria eradication (Joncour, 1959) and (ii) the massive use of organo-chlorine in agricultural settings during the 1950s (OMS, 1976). However, some cross-resistance exists between different groups of insecticides and emergence of resistance in vector populations is a major threat for the sustainability of malaria prevention through vector control in Africa (N'Guessan, 2009).

In West Africa, *A. gambiae* resistance to the four major classes of insecticides available for public health has been reported (Chandre et al., 1999; Elissa et al., 1993; Akogbeto and Yakoubou, 1999; Awolola et al., 2002; Fanello et al., 2003; Diabate et al., 2002). Pyrethroids are the only option for net treatment due to their relative safety for humans at low dosage, excito-repellent properties, rapid rate of knock-down and killing effects (Zaim et al., 2000). However, the success of the control methods is threatened by resistance to the main insecticides such as pyrethroids in malaria vectors.

Malaria vector resistance to insecticides in Benin is conferred by two main mechanisms: (1) alterations at site of action in the sodium channel via the *kdr* mutations and (2) an increase of detoxification and/or metabolism through high levels of multi-function oxidases (MFOs) and non-specific esterases (NSEs) (Corbel et al., 2007; Djogbénou et al., 2009; Djègbé et al., 2011; Aïzoun et al., 2013a).

A total of 48,819 Long-Lasting Insecticidal Nets, Permanet 2.0 (LLINs) were distributed to 47,524 households, with particular attention to children under-five and pregnant women, in October, 2008 and May, 2009 in the framework of President's Malaria Initiative of the U.S. Government in Oueme Department (Padonou et al., 2012). In addition, Beninese National Malaria Control Programme has implemented large-scale and free distribution of LLINs (OlysetNets) in July, 2011 throughout the entire country to increase coverage of LLINs. It is crucial that information on current status of *A. gambiae* s.l. permethrin, deltamethrin and DDT resistant populations should be investigated. This will properly inform control programs of the most suitable insecticides to use and facilitate the design of appropriate resistance management strategies.

Padonou et al. (2012) have shown that the main mechanism of resistance to pyrethroids is the mutation Leu1014F *kdr* allele in Seme district including Sekandji location. However, it would be useful to check if metabolic resistance conferred by detoxifying enzymes is also present in such *A. gambiae* populations. In fact, that will help to investigate multiple insecticide resistance mechanisms in *A. gambiae* Sekandji populations. In addition, Corbel et al. (2007) reported on multiple insecticide resistance mechanisms in *A. gambiae* Ladjì populations. These authors also mentioned that an experimental hut study carried out at Ladjì location in 2004 showed a rather low efficacy of permethrin treated nets at WHO recommended dosages against *A. gambiae* (Corbel et al., 2004) and this result underlines the need to investigate the role of enzymes in *A. gambiae* insecticide resistant populations through the use of classical synergists.

The main goal of this study was to explore the involvement of cytochrome P450 mono-oxygenases, esterases and glutathione S-transferases (GSTs) in permethrin, deltamethrin and DDT resistant *A. gambiae* s.l. populations from southern Benin by using classical synergists from 2008 to 2013.

METHODOLOGY

Study area

The study was carried out in the South of Benin, more precisely in Ladjì location, in the Cotonou district of Littoral Department and in Sekandji location, in the Seme district of Oueme Department (Figure 1). The choice of the study sites took into account the economic activities of populations, their usual protection practices against mosquito bites, the LLINs distribution recently in these localities and peasant practices to control farming pests. So, deltamethrin and permethrin were the two pyrethroid insecticides used in malaria vector control throughout LLINs distribution recently by Beninese National Malaria Control Programme. In addition, Permanets 2.0 were only distributed in Oueme Department whereas OlysetNets were distributed throughout the entire country. These factors have a direct impact on the development of insecticide resistance in the local mosquito vectors. Cotonou is characterized by a tropical coastal guinean climate with two rainy seasons (April to July and September to November). The mean annual rainfall is over 1,500 mm. Oueme has a climate with two rainy seasons (March to July and September to November). The temperature ranges from 25 to 30°C with the annual mean rainfall between 900 and 1,500 mm.

Mosquito sampling

A. gambiae s.l. mosquitoes were collected during the rainy seasons (March to July and September to November, 2008, 2010 and March to July, 2013) across Sekandji in the Seme district selected in south Benin. *A. gambiae* s.l. mosquitoes were also collected during the rainy seasons (April to July and September to November, 2008 and April to July, 2013) across Ladjì in the Cotonou district selected in south Benin. Larvae and pupae were collected on breeding sites using the dipping method. They were then kept in separated labeled bottles related to each locality. The samples were reared up to




Figure 1. Map of the study area.

adult emergence at the Centre de Recherche Entomologique de Cotonou, Benin (CREC) insectary. *A. gambiae* Kisumu, a reference susceptible strain was used as a control for the bioassay tests. Susceptibility tests were carried out following CDC protocols on unfed female mosquitoes aged 2 to 5 days old reared from larval and pupal collections. All susceptibility tests were conducted in the CREC laboratory at $25\pm 2^\circ\text{C}$ and 70 to 80% relative humidity.

CDC protocol

The principle of the CDC bottle bioassay is to determine the time it

takes an insecticide to penetrate an arthropod, traverse its intervening tissues, get to the target site, and act on that site relative to a susceptible control. Anything that prevents or delays the compound from achieving its objective of killing the arthropods contributes to resistance. Diagnostic doses that were applied in the current study were the doses recommended by CDC (Brogdon and Chan, 2010). These doses were checked on the *A. gambiae* Kisumu susceptible reference strain before being applied to field populations. For *A. gambiae s.l.*, the diagnostic dose of $12.5\ \mu\text{g}$ per bottle for deltamethrin and of $21.5\ \mu\text{g}$ per bottle for permethrin were used for the same diagnostic exposure time of 30 min whereas the diagnostic dose of $100\ \mu\text{g}$ per bottle for DDT was used for a




Figure 2. Diagram for performing the CDC bottle bioassay with synergists [CDC: Methods in *Anopheles* Research, 2010].

Figure 2: Diagram of performing the CDC bottle bioassay with synergists (CDC guideline, 2010)

diagnostic exposure time of 45 min. The choice of deltamethrin was justified by its use on Permanet 2.0 that was distributed free by the NMCP in October, 2008 and May, 2009 in Oueme Department, whereas permethrin is the insecticide used on OlysetNets that were distributed free by the NMCP in July, 2011 across the entire country. DDT was tested because of its intensive use in the past as well as to assess cross-resistance with permethrin and deltamethrin in localities surveyed.

The solutions were prepared and the bottles coated according to the CDC protocol (Brogdon and Chan, 2010). Fifteen to 20 unfed female mosquitoes aged 2 to 5 days old were introduced into four 250 ml Wheaton bottles coated with insecticide and one control bottle coated with acetone only. The number of dead or alive

mosquitoes was monitored at different time intervals (10, 20, 30, 40, 50, 60 min) in 2008 and (15, 30, 35, 40, 45, 60, 75, 90, 105, 120 min) in 2010 and 2013. This allowed us to determine the total percent mortality (Y axis) against time (X axis) for all replicates using a linear scale.

Biochemical assays using synergists

Synergists were used according to the protocol described by CDC (Brogdon and McAllister, 1998; Brogdon and Chan, 2010) following the procedure outlined in Figure 2. Samples that showed high resistance to permethrin in 2008 in Ladjì location from the Cotonou

district were exposed to the effects of two synergists: S.S.S-tributylphosphorotrithioate (DEF) (125 µg per bottle), which inhibits esterase activity and piperonyl butoxide (PBO) (400 µg per bottle), which inhibits oxidase activity. Piperonyl butoxide (PBO) is a pyrethroid synergist whereas S.S.S-tributylphosphorotrithioate is used in combination with PBO in order to explore multiple insecticide resistance mechanisms in *A. gambiae* Ladj populations. These two synergists were used separately and in combination. In a similar way, samples that showed high resistance to deltamethrin in 2010 in Sekandji location from the Seme district were exposed to the effects of these two same synergists (PBO and DEF). In addition, the samples that showed high resistance to DDT in 2013 in Sekandji locality were also exposed to the effects of the synergist: Ethacrynic acid (ETAA or EA) (80 µg per bottle), which inhibits glutathione S-transferases activity. This synergist was used in combination with DDT alone. Ethacrynic acid (ETAA or EA) is an organochlorine synergist such as dichlorodiphenyltrichloroethane (DDT). Approximately 125 mosquitoes were used for each synergist assay. The number of dead or alive mosquitoes was monitored at different time intervals (0, 10, 20, 30, 40, 50, 60 min) in 2008 and (0, 15, 30, 35, 40, 45, 60, 75, 90, 105, 120 min) in 2010 and in 2013. This test allowed us to compare the obtained percentages of dead mosquitoes (Y axis) against time (X axis) before the addition of the synergist (s) to those obtained after the addition of the synergist (s) (Figure 2).

Data analysis

The resistance status of mosquito samples was determined according to the CDC criteria (Brogdon and McAllister, 1998; Brogdon and Chan, 2010). The susceptibility thresholds at the diagnostic time of 30 min for pyrethroids and 45 min for organochlorines are: Mortality rate = 100%: the population is fully susceptible, Mortality rate < 100%: the population is considered resistant to the tested insecticides. Abbott's formula was not used in this study for the correction of mortality rates in the test-bottles because the mortality rates in all controls was always less than 5% (Abbott, 1987). To appreciate the effects of synergists PBO and DEF on *A. gambiae* Ladj and Sekandji permethrin and deltamethrin resistant populations in 2008 and in 2010, respectively and the effect of synergist ETAA on *A. gambiae* Sekandji DDT resistant populations in 2013, we used a Kruskal-Wallis test. The significance level was set at 5%. The software R-2.15.2. (R Development Core Team, 2011) was used for the statistical analysis.

RESULTS

Evolution of *A. gambiae* resistance to insecticides in Ladj and Sekandji locations from 2008 to 2013

The Kisumu strain (control) confirmed its susceptibility status as a reference strain. All female mosquitoes of *A. gambiae* Kisumu that were exposed to CDC bottles treated with permethrin 21.5 µg per bottle and DDT 100 µg per bottle in 2008 and in 2013, were dead and none of them could fly after 30 and 45 min, which represent the susceptibility threshold times or diagnostic times clearly defined by the CDC protocol. This confirmed that this strain was fully susceptible to these products in 2008 and 2013. In similar way, all female mosquitoes of *A. gambiae* Kisumu that were exposed to CDC bottles treated with

deltamethrin 12.5 µg per bottle in 2008, 2010 and in 2013 and with DDT 100 µg per bottle in 2008 and in 2013 were dead and none of them could fly after 30 and 45 min. This confirmed that this strain was fully susceptible to these products in 2008, 2010 and 2013.

A proportion of the *A. gambiae* Ladj population 28.56 and 11.67% in 2008 and 2013, respectively continued to fly again in the bottles following 30 min exposure to CDC bottles treated with permethrin. In addition, a large proportion of this population 91 and 97.68% in 2008 and 2013, respectively continued to fly again in the bottles following 45 min exposure to CDC bottles treated with DDT. This confirmed that *A. gambiae* Ladj population was highly resistant to these products (Table 1). No *A. gambiae* Ladj population was exposed to CDC bottles treated with permethrin 21.5 µg per bottle in 2010 (during our survey period) because it was difficult to collect a sufficient number of larvae and pupae of *A. gambiae* mosquitoes during this year in Ladj locality (Table 1).

A large proportion of the *A. gambiae* Sekandji populations (26%) continued to fly again in the bottles following 30 min exposure to CDC bottles treated with deltamethrin in 2010. This confirmed that these populations were resistant to this product (Table 1). Conversely to this resistance recorded in 2010, these populations were susceptible to the same product in 2008 and in 2013 when they had the same behavior facing deltamethrin as Kisumu susceptible reference strain. A large proportion of the *A. gambiae* Sekandji populations 85.92 and 97.15% in 2008 and 2013, respectively continued to fly again in the bottles following 45 min exposure to CDC bottles treated with DDT. This confirmed that these populations were highly resistant to this product (Table 1).

Effects of synergists PBO and DEF on *A. gambiae* Ladj populations resistant to permethrin in 2008

The data presented in Figure 3 show that after the addition of synergist PBO and DEF to permethrin 21.5 µg per bottle, the percentage of dead mosquitoes from Ladj is higher than the one obtained with permethrin alone. The use of either PBO or DEF synergist in bottles treated with permethrin 21.5 µg per bottle did not eliminate permethrin resistance, but significantly reduced the level with the mortality rate increasing from 71.25 to 84% ($p = 0.0053$) or from 71.25 to 80.80% ($p = 0.0425$), respectively. The use of the synergist combination DEF + PBO did not give the same result as the one obtained with PBO alone ($p = 0.0132$) or DEF alone ($p = 0.0017$). In addition, the use of synergist combination DEF + PBO did not restore the susceptibility in *A. gambiae* Ladj populations by rendering them susceptible to permethrin 21.5 µg per bottle as the reference strain Kisumu. These results suggest an implication of both mono-oxygenases and esterases in resistance of *A. gambiae* to pyrethroids.

Table 1. Mortality of *Anopheles gambiae* from the localities of Ladji and Sekandji after one and two hours exposure to CDC bottles treated with permethrin (21.5µg/bottle), deltamethrin (12.5µg/bottle), and DDT(100µg/bottle) from 2008 to 2013.

Locality	Year	Insecticide	Number tested	Mortality (%)	Resistance status
Kisumu (Ctrl)	2008	Permethrin	200	100	S
	2008	DDT	200	100	S
	2013	Permethrin	37	100	S
	2013	DDT	35	100	S
Ladji	2008	Permethrin	334	71.25	R
	2008	DDT	100	9	R
	2013	Permethrin	60	88.33	R
	2013	DDT	86	2.32	R
Kisumu(Ctrl)	2008	Deltamethrin	200	100	S
	2008	DDT	200	100	S
	2010	Deltamethrin	110	100	S
	2013	Deltamethrin	25	100	S
	2013	DDT	35	100	S
Sekandji	2008	Deltamethrin	86	100	S
	2008	DDT	72	14.08	R
	2010	Deltamethrin	47	74.07	R
	2013	Deltamethrin	41	100	S
	2013	DDT	70	2.85	R

Ctrl : Control

Effects of synergist ETAA on *A. gambiae* Ladji populations resistant to DDT

The analysis of Figure 4 shows that after the addition of synergist EA to DDT 100 µg/bottle, the percentage of dead mosquitoes from Ladji is slightly higher than the one obtained with DDT alone. The use of synergist EA in bottles treated with DDT 100 µg/bottle did not eliminate DDT resistance, and the mortality rate increased from 02.32 to 8.62% (P = 0.1179). These results show that GSTs may play a little role in *A. gambiae* Ladji resistance to DDT.

Effects of synergists PBO and DEF on *A. gambiae* Sekandji populations resistant to deltamethrin in 2010

The analysis of Figure 5 shows that after exposure to the synergist PBO prior to exposure to deltamethrin 1.25%, the percentage of dead mosquitoes from Sekandji on PBO was higher than that obtained with deltamethrin alone. The PBO synergist did not eliminate deltamethrin resistance but significantly reduced the level with the mortality rate increasing from 74.07 to 90.90% (p = 0.0122). The mortality rate recorded with deltamethrin + DEF was similar to the one obtained with deltamethrin alone (p = 0.5388). In addition, the use of the synergist

combination DEF + PBO did not give the same result as the one obtained with PBO alone (p = 02.99) or DEF alone (p = 0.0004). In addition, the use of synergist combination DEF + PBO did not restore the susceptibility in *A. gambiae* Sekandji populations by rendering them susceptible to deltamethrin 1.25% as the reference strain Kisumu. These results suggest an implication of mono-oxygenases in resistance of *A. gambiae* to pyrethroids.

Effect of synergist ETAA or EA on *Anopheles gambiae* Sekandji populations resistant to DDT in 2013

The analysis of Figure 6 shows that after the addition of synergist ETAA to DDT 100 µg per bottle, the percentage of dead mosquitoes from Sekandji is lower than the one obtained with DDT alone. The use of synergist ETAA in bottles treated with DDT 100 µg per bottle did not eliminate DDT resistance, and the mortality rate decreased from 2.85 to 0% (P = 0.5114). These results show that GSTs may play no role in *A. gambiae* Sekandji resistance to DDT.

DISCUSSION

A. gambiae Ladji populations were resistant to permethrin




Figure 3. Effects of synergists PBO and DEF on *Anopheles gambiae* Ladji populations resistant to permethrin in 2008.




Figure 4. Effects of synergist ETAA on *Anopheles gambiae* Ladji populations resistant to DDT.




Figure 5. Effects of synergists PBO and DEF on *Anopheles gambiae* Sekandji populations resistant to deltamethrin in 2010.




Figure 6. Effect of synergist ETAA or EA on *Anopheles gambiae* Sekandji populations resistant to DDT in 2013.

in 2008 and still remained resistant to the same product in 2013. The pyrethroid resistance observed in *A. gambiae* Ladji populations may be due to the presence of several environmental pollutants and pesticide residues from the neighbouring peri-urban cities and farms to the coastal location of Ladji, a peri-urban locality located in the town of cotonou. This locality is crossed by the Nokoue Lake's streams. These streams sweep and converge these environmental pollutants and pesticide residues in Ladji locality. These xenobiotics available in larval breeding sites in Ladji may be one of the possible factors selecting for pyrethroid resistance in *A. gambiae* populations in this locality. A similar pattern was already observed with *A. gambiae* Agbalilamè permethrin resistant populations (Aïzoun et al., 2013a). In the current study, in 2008, metabolic resistance conferred by detoxifying enzymes such as cytochrome P450 mono-oxygenases and esterases were found to play a role in *A. gambiae* Ladji permethrin resistant populations.

In addition, after the addition of both synergists PBO and DEF on *A. gambiae* Ladji permethrin resistant populations, we have not still obtained fully susceptibility. These results showed that there were other resistance mechanisms which were not synergizable by PBO and DEF. In southern Benin, Corbel et al. (2007) have already reported on multiple insecticide resistance mechanisms in *A. gambiae* Ladji populations. Among these mechanisms, there were mixed function oxidase (MFO) and α -esterase with the presence of *Kdr* at high frequency (80%). However, even if the Leu-Phe *kdr* mutation is the most important resistance mechanism in these *A. gambiae* Ladji populations, metabolic resistance conferred by detoxifying enzymes is also an indication of phenotypic resistance to permethrin.

A. gambiae Ladji populations were resistant to DDT in 2008 and still remained resistant to the same product in 2013 with an increase in the resistance level. According to Akogbeto and Yakoubou (1999), the emergence of DDT resistance recorded in *A. gambiae* from meridian regions was related to two phenomena: the massive use of DDT and dieldrin for house-spraying applications in southern villages from 1953 to 1960 during WHO programmes of malaria eradication and the massive use of organochlorine in agricultural settings during the 1950s (OMS, 1976). In addition, previous studies conducted by Akogbeto et al. (2006) and Corbel et al. (2007) have already showed a cross-resistance to pyrethroid and DDT in *A. gambiae* Ladji populations. A recent study conducted in 2013 showed that GSTs may play a little role in *A. gambiae* Ladji DDT resistant populations using CDC biochemical assays (Aïzoun et al., 2013b). This result could explain in part the other resistance mechanisms involved in *A. gambiae* Ladji permethrin resistant populations because this permethrin resistance was not synergizable by PBO and DEF in 2008. This result also shows that the three categories of enzymes namely esterases, cytochrome P450 mono-oxygenases

and glutathione S-transferases which were typically involved in insecticide resistance in malaria vectors were all present in *A. gambiae* Ladji populations.

A. gambiae Sekandji populations were susceptible to deltamethrin in 2008 and 2013. But in 2010, these populations were resistant to this product. According to Padonou et al. (2012), the deltamethrin resistance recorded in *A. gambiae* Seme populations in 2010 was not due to PermaNet2.0 distribution by National Malaria Control Programme (NMCP), and the high frequency of the mutation L1014F *kdr* allele could explain this resistance. These authors also mentioned that the selective pressure exerted by the promotion of mosquito nets by the Health Ministry and the free distribution of LLINs in Oueme region, causing the *kdr* increase within *A. gambiae* populations is doubtful. The *kdr* frequency recorded in Seme district including Sekandji locality by Padonou et al. (2012) in 2010 was 0.87. During the same year, metabolic resistance conferred by cytochrome P450s was detected in *A. gambiae* Sekandji deltamethrin resistant populations in the current study. A recent study carried out by Aïzoun et al. (2013a) in Seme district, precisely in Agbalilame locality also suggested an implication of mono-oxygenases in resistance of *A. gambiae* s.l. to pyrethroids. Indeed, Padonou et al. (2012) showed that the main mechanism of pyrethroid resistance in *A. gambiae* Seme was L1014F *kdr* mutation. As it was observed with *A. gambiae* Ladji populations, even if the Leu-Phe *kdr* mutation is the most important resistance mechanism in *A. gambiae* Sekandji populations collected in Seme district, metabolic resistance conferred by detoxifying enzymes is also an indication of phenotypic resistance to deltamethrin.

A. gambiae Sekandji populations were resistant to DDT in 2008 and still remained resistant to the same product in 2013 with an increase in the resistance level. The cause of this resistance level increasing was the same as the one observed with *A. gambiae* Ladji DDT resistant populations. In fact, Sekandji is also located in the meridian region like Ladji. In the current study, after the addition of synergist ETAA on *A. gambiae* Sekandji DDT resistant populations, the mortality rate decreased and Glutathione S-transferases therefore may play no role in these *A. gambiae* Sekandji DDT resistant populations. A similar pattern was already observed with *A. gambiae* Bohicon and Parakou populations (Aïzoun et al., 2013b). In some cases, the use of synergists at the same time as the application of insecticide could inhibit the penetration of the insecticide through the cuticle, therefore reducing the amount of insecticide entering the insect's body (Martin et al., 1997), the result of which was that the toxicity effect would also be reduced.

Conclusion

This study shows that *A. gambiae* s.l. populations from southern Benin were resistant to both DDT and pyrethroids.

The resistance level of these populations to these products has varied or changed in an interval of five years. Even if the Leu-Phe *kdr* mutation is the most important resistance mechanism in these *A. gambiae s.l.* populations, metabolic resistance conferred by detoxifying enzymes is also an indication of phenotypic resistance to both DDT and pyrethroids in southern Benin.

REFERENCES

- Abbott WS (1987). A method of computing the effectiveness of an insecticide. *J. Am. Mosq. Control Assoc.* 3(2):302–303.
- Aïzoun N, Aïkpon R, Padonou GG, Oussou O, Oké-Agbo F, Gnanguenon V, Ossè R, Akogbéto M (2013a). Mixed-function oxidases and esterases associated with permethrin, deltamethrin and bendiocarb resistance in *Anopheles gambiae s.l.* in the south-north transect Benin, West Africa. *Parasit. Vectors* 6:223.
- Aïzoun N, Aïkpon R, Agossa F, Gnanguenon V, Padonou GG, Akogbéto M (2013b). Insecticide resistance and effect of synergist Ethacrynic acid (ETAA or EA) on DDT resistance cross to permethrin in *Anopheles gambiae s.l.* in the south-north transect Benin, West Africa. *J. Parasitol. Vector Biol.* (in Press).
- Akogbeto M, Djouaka R, Kinde-Gazard DA (2006). Screening of pesticide residues in soil and water samples from agricultural settings. *Malar. J.* 5:22.
- Akogbeto M, Yakoubou S (1999). Résistance des vecteurs du paludisme vis-à-vis des pyrèthrinoides utilisés pour l'imprégnation des moustiquaires au Bénin, Afrique de l'Ouest. *Bull. Soc. Pathol. Exot.* 92:123-130.
- Awolola TS, Brooke BD, Hunt RH, Coetze M (2002). Resistance of the malaria vector *Anopheles gambiae s.s.* to pyrethroid insecticides, in south-western Nigeria. *Ann. Trop. Med. Parasitol.* 96(8):849-852.
- Brogdon WG, McAllister JC (1998). Simplification of adult mosquito bioassays through use of time-mortality determinations in glass bottles. *J. Am. Mosq. Control. Assoc.* 14(2):159–64.
- Brogdon W, Chan A (2010). Guidelines for Evaluating Insecticide Resistance in Vectors using the CDC Bottle Bioassay/Methods in anopheles research. Second edition. CDC Atlanta USA: CDC technical report 343.
- Chandre F, Darriet F, Manga L, Akogbeto M, Faye O, Mouchet J, Guillet P (1999). Status of pyrethroid resistance in *Anopheles gambiae sensu lato*. *Bull. WHO.* 77(3):230-234.
- Corbel V, Chandre F, Brengues C, Akogbeto M, Lardeux F, Hougard JM, Guillet P (2004). Dosage dependent effects of permethrin-treated nets on the behaviour of *Anopheles gambiae* and the selection of pyrethroid resistance. *Malar. J.* 3:22.
- Corbel V, N'Guessan R, Brengues C, Chandre F, Djogbenou L, Martin T, Akogbeto M, Hougard JM, Rowland M (2007). Multiple insecticide resistance mechanisms in *Anopheles gambiae* and *Culex quinquefasciatus* from Benin, West Africa. *Acta. Trop.* 101:207-216.
- Coz J, Davidson G, Chauvet G, Hamon J (1968). La résistance des anophèles aux insecticides en Afrique Tropicale et à Madagascar. *Cah ORSTOM, Sér Ent Méd Parasitol.* 38:14.
- Diabaté A (1999). Evaluation de la résistance des vecteurs du paludisme vis-à-vis des pyrèthrinoides au Burkina Faso. Thèse de 3^e cycle, Université de Ouagadougou. p 128.
- Diabate A, Baldet T, Chandre F, Guiguemde RT, Brengues C, Guillet P, Hemingway J, Hougard JM (2002). First report of the *kdr* mutation in *Anopheles gambiae* M form from Burkina Faso, West Africa. *Parassitologia* 44(3-4):157-158.
- Djègbé I, Boussari O, Sidick A, Martin T, Ranson H, Chandre F, Akogbéto M, Corbel V (2011). Dynamics of insecticide resistance in malaria vectors in Benin: first evidence of the presence of L1014S *kdr* mutation in *Anopheles gambiae* from West Africa. *Malar. J.* 10:261.
- Djogbéno L, Pasteur N, Akogbéto M, Weill M, Chandre F (2009). Insecticide resistance in the *Anopheles gambiae* complex in Benin: a nationwide survey. *Med. Vet. Entomol.* 69:160-164.
- Elissa N, Mouchet J, Riviere F, Meunier JY, Yao K (1993). Resistance of *Anopheles gambiae s.s.* to pyrethroids in Cote d'Ivoire. *Ann. Soc. Belge. Méd. Trop.* 73(4):291-294.
- Fanello C, Petrarca V, della Torre A, Santolamazza F, Dolo G, Coulibaly M, Allouche A, Curtis CF, Toure YT, Coluzzi M (2003). The pyrethroid knock-down resistance gene in the *Anopheles gambiae* complex in Mali and further indication of incipient speciation within *A. gambiae s.s.* *Insect. Mol. Biol.* 12(3):241-245.
- Joncour G (1959). Lutte anti-palustre au Dahomey. Rapport n° 13, Ministère de la Santé Publique. p 10.
- Martin SH, Ottea JA, Leonard BR, Graves JB, Burris E, Micinski S, Church GE (1997). Effects of selected synergists and insecticide toxicity in tobacco budworms (Lepidoptera: Noctuidae) in laboratory and field studies. *J. Econ. Entomol.* (3):723-721.
- N'Guessan RK (2009). Insecticide resistance in the West African malaria vector *Anopheles gambiae* and investigation of alternative tools for its delay. PhD Thesis, Wageningen University, Netherlands. p 222.
- OMS (1976). Résistance des vecteurs et des réservoirs de maladies au pesticides 22 me rapport du comité OMS d'experts des insecticides. *Cah. Tech. Afro.* 585:30-31.
- Padonou GG, Sezonlin M, Ossè R, Aizoun N, Oké-Agbo F, Oussou O, Gbédjissi G, Akogbéto M (2012). Impact of three years of large scale Indoor Residual Spraying (IRS) and Insecticide Treated Nets (ITNs) interventions on insecticide resistance in *Anopheles gambiae s.l.* in Benin. *Parasit. Vectors* 5:72.
- Zaim M, Aitio A, Nakashima N (2000). Safety of pyrethroid-treated mosquito nets. *Med. Vet. Entomol.* 14:1-5.